## Evidence-Based Medicine

## for Primary Care and Internal Medicine



Editors: Brian Haynes & Paul Glasziou

## **Contents**

orboso and brotogoro		manticution of intervenience and as militial balancia and the many and	
BM notebook		motivational interviews, and cognitive behavioural therapy red excessive alcohol consumption in primary care	179
ottings	162	An intensive lifestyle intervention reduced weight and	1/3
ife after the EBM workshop	163	cardiovascular disease risk factors in obese participants with ty	/ne
mart health choice essentials	164	2 diabetes at 1 year in an interim analysis	180
etter	166	Irrigation of simple lacerations with tap water or sterile saline	in
		the emergency department did not differ for wound infections	181
vidently	167	<b>N</b>	
		<b>Diagnosis</b>	
herapeutics leview: vaccines against human papillomavirus prevent cervical		Review: some screening tests for dementia in older people are	101
ancer precursors in young women for at least 3 years	168	accurate and practical for use in primary care Review: anti—cyclic citrullinated peptide antibody is a more	182
leview: rosiglitazone increases risk of MI but does not differ from	100	specific test for rheumatoid arthritis than rheumatoid factor	183
ther drugs for CV death in type 2 diabetes	169	specific 1031 for modification difficults must modificate tactor	.00
osiglitazone increased heart failure but did not differ from		Aetiology	
netformin plus sulphonylurea for other CV outcomes at interim		Review: increased waist circumference or waist-to-hip ratio is	
nalysis	170	associated with increased risk of cardiovascular events	184
eview: regular brisk walking improves cardiovascular risk factors	171		
n healthy sedentary adults Variaria radused major strake more than genicia in elderly	171	Quality improvement	J
Varfarin reduced major stroke more than aspirin in elderly atients with atrial fibrillation in primary care	172	Reminders in echocardiography reports increased use of $\beta$ blo	ckers 1 <b>85</b>
lymptomatic use of beclomethasone-albuterol and regular use of	1/2	in reduced left ventricular ejection fraction	103
eclomethasone did not differ for control of mild asthma	173	Clinical prediction guide	
nduction of labour post-term or expectant management with		Review: evidence from single studies shows that a few fall risl	k
erial fetal monitoring had similar neonatal outcomes	174	assessment tools can predict falls in elderly people	186
rednisolone plus paracetamol (acetaminophen) was as effective		The ankle brachial index independently predicted allcause mor	tality
s indomethacin plus paracetamol but had fewer adverse effects	175	in high-risk patients with peripheral arterial disease	187
acute gout-like arthritis	175		100 100
ntranasal fentanyl and intravenous morphine did not differ for ain relief in children with closed long-bone fractures	176	Resource review	188, 189
leview: on-demand maintenance therapy with proton pump	170	Additional articles abstracted in ACP Journal Club	190
shibitors is as effective as continuous therapy for non-erosive		Additional diffices abstracted in ACI Journal Clob	170
ORD	177	Other articles noted	191
leaning of the perineal/genital area before urine collection from			
pilet-trained children prévented sample contamination	178	<b>Glossary</b> Inside	e back cov