


# Evidence-Based Medicine



**Editors:** Brian Haynes & Paul Glasziou

## Contents

Purpose and procedure	2		
Additional articles abstracted in <i>ACP Journal Club</i>	3		
EBM notebook: "Double blind, you are the weakest link — goodbye!"	4		
Resource corner	7		
<b>Therapeutics</b>			
Review: varicella vaccination is effective in children	9		
Oseltamivir was safe and effective for prophylaxis of influenza in the frail elderly	10		
Review: placebo is better than no treatment for subjective continuous outcomes and for treatment of pain	11		
Omaliuzumab reduced inhaled corticosteroid use and exacerbations in childhood allergic asthma	12		
Review: angiotensin-converting enzyme inhibitors reduce the progression of non-diabetic renal disease	13		
Predischarge coronary angiography was better than exercise testing for reducing hospital use after low-risk chest pain	14		
Carvedilol reduced mortality and morbidity caused by myocardial infarction in patients with left ventricular dysfunction	15		
Long term and short term oral anticoagulation treatments were equivalent for venous thromboembolism	16		
3 months and 1 year of oral anticoagulant treatment were equivalent for idiopathic proximal deep venous thrombosis	17		
Home monitoring service was more effective than usual care in patients with essential hypertension	18		
Review: antibiotics modestly increase remission rates in acute bacterial conjunctivitis	19		
Dietary fibre supplementation with psyllium or gum arabic reduced faecal incontinence in community-living adults	20		
		17β-estradiol reduced depressive and somatic symptoms in perimenopausal women	21
		Brief psychodynamic interpersonal treatment after deliberate self poisoning reduced suicidal ideation and deliberate self harm	22
		Support by another mother and a child-life specialist decreased anxiety in mothers of children with chronic illnesses	23
		Review: cannabinoids and codeine have similar effects on pain relief, but cannabinoids commonly cause psychotropic adverse effects	24
		Review: cannabinoids control chemotherapy-induced nausea and vomiting but increase the risk of side effects	25
		Surgery relieved symptoms but decreased survival more than medical treatment in gastro-oesophageal reflux disease	26
		<b>Prognosis</b>	
		Progression of disease in HIV infected children slowed after the first year of life	27
		More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery	28
		<b>Etiology</b>	
		Cardiovascular events were increased in patients with high cardiovascular risk who had albuminuria	29
		<b>Quality improvement</b>	
		Recall to a general practitioner or to a nurse clinic improved assessment in patients with coronary artery disease	30
		<b>Clinical prediction guide</b>	
		The Framingham scores overestimated the risk for coronary heart disease in Japanese, Hispanic, and native American cohorts	31
		A clinical prediction model did well in diagnosing paediatric group A β-haemolytic streptococcal pharyngitis	32