Review: observational studies adjusting for socioeconomic status and lifestyle show no association between HRT and CAD

Humphrey LL, Chan BK, Sox HC. Postmenopausal hormone replacement therapy and the primary prevention of cardiovascular disease. *Ann Intern Med* 2002;137:273–84.

QUESTION: What is the effectiveness of postmenopausal hormone replacement therapy (HRT) for primary prevention of cardiovascular disease?

Data sources

Studies were identified by searching Medline and Cochrane databases from 1966 to December 2000 and by reviewing bibliographies of relevant studies and other publications.

Study selection

Studies were selected if they were randomised controlled trials (RCTs), cohort studies, or case control studies that assessed the use of HRT for primary prevention of cardiovascular disease in postmenopausal women and if English language abstracts were available.

Data extraction

Data were extracted on study design, type of HRT (unopposed oestrogen or oestrogen plus progestin),

definitions of cardiovascular disease, and potential risk factors included in multivariate models. 2 investigators independently assessed the quality of individual studies. Formal review and meta-analysis were limited to studies of good or fair quality and included RCTs, cohort studies with internal controls, and population based case control studies with ≥ 3 years of follow up. HRT use was classified as current, past, ever, or any (combined current, past, or ever).

Main results

Only the results for coronary artery disease (CAD) incidence are reviewed here (3 cohort studies, 9 case control studies, and 1 small RCT). Studies that did not adjust for socioeconomic status (SES) found that current and past use of HRT reduced CAD incidence, whereas studies that adjusted for SES found no association between any measure of HRT use and CAD (table). Similar results were found when analyses were stratified by studies that adjusted for alcohol consumption, exercise, or both.

Conclusions

Meta-analysis of studies that adjust for socioeconomic status or alcohol consumption and exercise show that current, past, ever, or any use of HRT does not reduce coronary artery disease (CAD). Studies that do not adjust for these factors show a reduced risk for CAD with current and past use.

Source of funding:
Agency for Healthcare
Research and Quality.

For correspondence:
Dr L L Humphrey,
Veterans Affairs
Medical Center and
Oregon Health and
Science University,
Portland, OR, USA.
humphreyl@ohsu.edu

Abstract and
commentary also
appear in *ACP Journal
Club*.

Relative risk for coronary artery disease according to use of hormone replacement therapy (HRT)

Use of HRT	Relative risk (95% credible interval)	
	Unadjusted	Adjusted†
Current	0.71 (0.64 to 0.78)	0.97 (0.82 to 1.16)‡
Past	0.78 (0.69 to 0.87)	1.07 (0.90 to 1.27)‡
Ever	Not reported	1.11 (0.84 to 1.53)‡
Any	Not reported	1.04 (0.79 to 1.44)‡

*Includes data from studies that did not adjust for socioeconomic status.

†Includes data from studies that adjusted for socioeconomic status.

‡Not statistically significant.

COMMENTARY

The 2 meta-analyses by Humphrey *et al* and Nelson *et al*, and much of the medical literature, do not clearly define postmenopausal HRT. Although authors often claim that many observational studies show the benefits of HRT on CAD risk, the data almost completely relate to unopposed oestrogen.¹ Few epidemiological studies have evaluated oestrogen plus progestin administered as continuous combined therapy. A limited number of studies were included in the meta-analyses by Humphrey *et al* and Nelson *et al*, and the included studies sometimes failed to specify type of HRT. Studies published in the 1980s and earlier probably refer to unopposed oestrogen therapy.

The lack of specificity in type of HRT is problematic because of the potential for different effects of oestrogen alone and oestrogen plus progestin. It is important to note that the meta-analyses used a common protocol to apply standard criteria for quality² and included only studies of higher quality. For studies assessing CAD outcomes, the authors identified a relation between poorer quality studies and greater protection against CAD.

Changing patterns of use (oestrogen alone replaced by combined oestrogen plus progestin) and indications for use (eg, starting women on treatment before menopause to reduce bone loss) may account, in part, for differences in results. Also, acute effects of combined treatment (eg, the prothrombotic and proinflammatory effects of progestins) may be missed in prospective studies because of lack of attention to early events and follow up.

Data from both epidemiological studies and RCTs of HRT consistently show other thrombotic effects, such as stroke and pulmonary embolism. This argues, at least in part, against a bias related to this pathway in the observational studies. Although socioeconomic status (SES) appears as one other important variable in this analysis, the authors note that the range of cardiovascular risk factors controlled for in observational studies also varied substantially. The authors, however, did not point out that studies that controlled for SES observed similar results before and after such control, which suggests that this is not an explanation for the discrepancy between RCT and observational study results. Is confounding by indication changing over time concurrent with changing patterns of drug combination? Is SES merely a marker for more recent studies that evaluate oestrogen plus progestin? Alternatively, is the timing of exposure in relation to menopause the explanatory factor? Animal studies suggest that oestrogens have beneficial effects in the early stages of atherogenesis, but reduced beneficial effects in the final stages of plaque complications.³

continued on next page

Review: risks and benefits of HRT comparing various sources of evidence

Nelson HD, Humphrey LL, Nygren P, et al. Postmenopausal hormone replacement therapy: scientific review. *JAMA* 2002;288:872–81.

QUESTION: What are the benefits and harms of hormone replacement therapy (HRT) for primary prevention of cardiovascular disease, thromboembolism, osteoporosis, cancer, dementia, and cholecystitis?

Data sources

Studies were identified by searching Medline (1966–2001), HealthSTAR (1975–2001), and the Cochrane Controlled Trials Register; reviewing bibliographies of relevant studies, reviews, and editorials; and contacting experts.

Study selection

Studies were selected if they included a comparison group of HRT non-users and reported data relating to HRT use and clinical outcomes of interest. Studies were excluded if the sample was selected according to previous events or conditions associated with higher risks of targeted outcomes.

Data extraction

Data were extracted on study design and type of HRT (unopposed oestrogen or oestrogen plus progestin). 2 reviewers independently assessed study quality as good, fair, or poor using the US Preventive Services Task Force criteria.

Main results

The findings of the meta-analyses (relative risks) for some outcome categories are summarised in the table with corresponding hazard ratios from the recent Women's Health Initiative (WHI). The results of the meta-analysis showed that HRT reduced the risk of wrist fractures, vertebral fractures, colon cancer, and dementia, and increased the risk of stroke, thromboembolic events, breast cancer, and cholecystitis (table). Findings

for some of these outcomes were available from the WHI randomised controlled trial and differed (in terms of statistical significance when using adjusted results for secondary outcomes) for vertebral fractures, colon cancer, coronary heart disease events, and stroke (table).

Conclusions

Results of this meta-analysis of primarily observational studies and those of a large randomised controlled trial (Women's Health Initiative [WHI]) both show that hormone replacement therapy (HRT) increases risk of thromboembolic events. The meta-analysis shows no effect of HRT on coronary heart disease events, whereas the WHI found an increased risk.

Sources of funding:
Agency for Healthcare
Research and Quality
and Portland Veterans
Affairs Medical Center
Women's Health
Fellowship.

For correspondence:
Dr H D Nelson, Oregon
Health and Science
University, Portland,
OR, USA.
nelsonh@ohsu.edu

Abstract and
commentary also
appear in *ACP Journal
Club*.

Risks associated with ever use of HRT from Nelson et al and from the Women's Health Initiative [WHI]*

Outcomes	Relative risk (95% CI) from Nelson et al	Hazard ratio (CI) from WHI
Hip fractures	0.76 (0.56 to 1.01)	0.66 (0.33 to 1.33)‡
Vertebral fractures	0.60 (0.36 to 0.99)†	0.66 (0.32 to 1.34)‡
Colon cancer	0.80 (0.74 to 0.86)†	0.63 (0.32 to 1.24)‡
Coronary heart disease events	0.91 (0.67 to 1.33)	1.29 (1.02 to 1.63)†
Stroke	1.12 (1.01 to 1.23)†	1.41 (0.86 to 2.31)‡
Thromboembolic events (current use)	2.14 (1.64 to 2.81)†	2.11 (1.26 to 3.55)†‡
Breast cancer (≥5 y HRT)	1.23 to 1.35	1.26 (1.00 to 1.59)†

*Abbreviations defined in glossary; meta-analyses based on a random effects model.

†Statistically significant.

‡Adjusted CIs.

COMMENTARY—continued from previous page

How can drug formulation change over time and most associations observed in epidemiological studies (ie, breast cancer, pulmonary embolus, stroke, colon cancer, cholecystitis, and osteoporotic fracture) be consistent, and yet CAD outcomes diverge? Consistent evidence across the RCTs argues against chance.

The consistency between non-RCTs and RCTs for non-cardiac outcomes is reassuring and supports other evaluations of the contribution of different study designs to evaluation of medical therapies.^{1–7} Such studies comparing designs may be limited in power, but on average, well designed observational studies show a similar magnitude of estimated benefits as RCTs. We should not focus solely on study design but must also consider the formulation and timing of use of postmenopausal HRT that is being evaluated.

Graham Colditz, MD
Channing Laboratory
Boston, Massachusetts, USA

- Stampfer MJ, Colditz GA. Estrogen replacement therapy and coronary heart disease: a quantitative assessment of the epidemiologic evidence. *Prev Med* 1991;20:47–63.
- Harris RP, Helfand M, Woolf SH, et al. Current methods of the US Preventive Services Task Force: a review of the process. *Am J Prev Med* 2001;20:21–35.
- Mikkola TS, Clarkson TB. Estrogen replacement therapy, atherosclerosis, and vascular function. *Cardiovasc Res* 2002;53:605–19.
- Colditz GA, Miller JN, Mosteller F. How study design affects outcomes in comparisons of therapy. I: Medical. *Stat Med* 1989;8:441–54.
- Ioannidis JP, Haidich AB, Pappa M, et al. Comparison of evidence of treatment effects in randomized and nonrandomized studies. *JAMA* 2001;286:821–30.
- Benson K, Hartz AJ. A comparison of observational studies and randomized, controlled trials. *N Engl J Med* 2000;342:1878–86.
- Concato J, Shah N, Horwitz RI. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000;342:1887–92.