


Evidence-Based Medicine

Editors: Brian Haynes & Paul Glasziou



EBM notebook

Using Bayes' nomogram to help interpret odds ratios 132

Resource review

Therapeutics

Review: continuing treatment with antidepressants reduces the rate of relapse or recurrence of depressive symptoms regardless of duration of treatment before or after randomisation 137

Review: electroconvulsive therapy reduces depressive symptoms 138

Review: restricted dietary salt intake can reduce blood pressure but does not reduce death or cardiovascular morbidity 139

Rhythm control strategies were not better than rate control strategies for atrial fibrillation 140

Rate control was not inferior to rhythm control for recurrent persistent atrial fibrillation 141

An intensive intervention reduced cardiovascular and microvascular events in type 2 diabetes and microalbuminuria 142

Combination ACE inhibitor and angiotensin receptor blocker therapy was better than monotherapy in non-diabetic renal disease 143

ACE inhibitors reduced cardiovascular events and all cause mortality in elderly people with hypertension 144

Review: long acting β_2 agonists are better than short acting β_2 agonists in chronic asthma 145

Adjunctive treatment with eplerenone reduced morbidity and mortality in acute myocardial infarction 146

Review: some non-surgical interventions improve symptoms in carpal tunnel syndrome 147

Mitoxantrone slowed progression of disability and reduced relapses in multiple sclerosis 148

A collaborative care management programme in a primary care setting was effective for older adults with late life depression 149

Buprenorphine improved treatment retention in patients with heroin dependence 150

Subcutaneous and intravenous infusions were similarly well accepted, safe, and feasible in older patients with dehydration 151

Morphine sulphate analgesia did not affect diagnostic accuracy in undifferentiated abdominal pain 152

Review: herbal medicinal products seem to be effective and safe in nonulcer dyspepsia 153

A topical cream containing glucosamine and chondroitin sulphate reduced joint pain in osteoarthritis of the knee 154

Diagnosis

The WHO-5 Wellbeing index performed the best in screening for depression in primary care 155

Review: fair evidence supports several tests in investigating fever of unknown origin 156

Aetiology

Diastolic dysfunction was predictive of all cause mortality 157

Quality improvement

The benefits of nurse led secondary prevention clinics continued after 4 years 158

Economics

Coxibs were not cost effective for arthritis pain in patients with average risk of ulcer complications 159

Psychotherapy or paroxetine did not reduce abdominal pain, but may improve quality of life in irritable bowel syndrome 160