

Therapeutics

Review: a home-like birth environment has beneficial effects on labour and delivery

Hodnett ED. *Home-like versus conventional institutional settings for birth*. *Cochrane Database Syst Rev* 2002;(1):CD000012 (latest version 13 Jul 2001).

QUESTION: In women at low risk for obstetric complications, what are the effects of a home-like birth environment compared with those of a conventional labour ward on labour and birth outcomes?

Source of funding: no external funding.

For correspondence: Professor E D Hodnett, University of Toronto, Toronto, Ontario, Canada. ellen.hodnett@utoronto.ca

Data sources

Studies were identified by using the search strategy developed for the Cochrane Pregnancy and Childbirth Group, which included searching Medline and the Cochrane Controlled Trials Register, and by hand-searching 38 relevant journal titles.

Study selection

Studies were selected if they were randomised controlled trials (RCTs) or quasi-RCTs that compared a home-like institutional birth setting with a conventional institutional birth setting in low risk pregnant women.

Data extraction

Data were extracted on methodological quality, including allocation concealment procedure, patient characteristics, birthing centre characteristics, and outcomes. Outcomes were intrapartum medical interventions, intrapartum and postpartum complications, type of delivery, perinatal mortality, feeding practices and problems, neonatal health outcomes, and adjustment to mothering.

Main results

5 RCTs and 1 quasi-RCT (n=8677) were included. 19% to 77% of women allocated to a home-like birth setting were transferred to conventional care before or during labour. Women who gave birth in the home-like setting had less use of pain medication during labour, were less likely to be immobile during labour, had fewer fetal heart rate abnormalities, had fewer operative deliveries, and were less dissatisfied with their care than were women in the conventional labour ward setting (table). Groups did not differ for discontinuation of breast feeding at 8 weeks (table). A trend toward higher perinatal mortality was seen in the home-like setting [p=0.066]* (table).

Conclusion

In women at low risk for obstetric complications, giving birth in a home-like birth environment is associated with

less analgesia use, operative delivery, and fetal abnormalities and greater satisfaction with care.

*p Value calculated from data in article.

COMMENTARY

Home-like birth environments, in which low risk women are looked after mainly by professional midwives, have shown such benefits as greater maternal satisfaction¹ and decreases augmentation, use of analgesia, operative deliveries, and fetal heart rate abnormalities.

In the review by Hodnett, the results show that when women are looked after in home-like birth environments they are more likely to be mobile during the first stage of labour and more likely to deliver in an upright or semirecumbent position. These 2 factors may contribute to the substantial decrease in augmentation rates, use of analgesia, operative deliveries, and abnormal fetal heart rate tracings.

The interventions in this review can be applied to most developed and developing countries. Analgesia (epidural and spinal) or operative delivery are an enormous cost burden to health service providers, and it has been shown that the average uncomplicated vaginal birth costs 68% less in a home setting than in a hospital.²

The concern is that the delay in operative deliveries or delay of interventions may contribute to an increase in perinatal mortality. Although a trend existed toward increased perinatal mortality, the incidence was low and within normal limits.³ It is important that healthcare providers giving care to women in home-like birth environments continue to observe the women during labour and birth so that fetal compromise is recognised early and dealt with effectively.

The use of home-like birth environments for low risk women has beneficial effects on parturients. The results of the review by Hodnett support the goals of the Better Births Initiative.⁴ Financial shortages are major problems in developing countries. Introducing home-like birth environments with good clinical care could be a cost-effective healthcare alternative but warrants further attention, specifically to perinatal mortality.

Cheryl Nikodem, DCur
Tembisa Hospital Effective Care Research Unit
Midrand
South Africa

1 Longworth L, Ratcliffe J, Boulton M. *Health Soc Care Community* 2001;9:404–13.

2 Anderson RE, Anderson DA. *J Nurse Midwifery* 1999;44:30–5.

3 Gilbert RE, Tookey PA. *BMJ* 1999;319:483–7.

4 Effective Health Care Alliance Programme. Better births initiative (www.liv.ac.uk/1stm/bbmainpage.html).

Home-like birth setting v conventional labour ward†

Outcomes	Number of trials	Weighted event rates			
		Home-like setting	Conventional setting	RRR (95% CI)	NNT (CI)
Any intrapartum analgesia	4	69%	75%	8% (5 to 10)	18 (13 to 27)
Immobility during labour	1	31%	39%	19% (10 to 27)	14 (10 to 28)
FHR abnormality	2	19%	25%	23% (15 to 30)	18 (13 to 29)
Operative delivery	5	15.2%	17.4%	12% (3 to 20)	46 (27 to 200)
Dissatisfaction with care	2	27%	70%	62% (50 to 71)	3 (2 to 3)
Stopped breast feeding within 6 to 8 weeks	2	1.26%	1.26%	1% (–60 to 149)	Not significant
				RRI (CI)	NNH
Perinatal mortality	4	0.68%	0.38%	82% (–3 to 243)	Not significant

†FHR = fetal heart rate. Other abbreviations defined in glossary; RRR, RRI, NNT, NNH, and CI calculated from data in article using fixed effects.



Review: a home-like birth environment has beneficial effects on labour and delivery

Evid Based Med 2002 7: 105
doi: 10.1136/ebm.7.4.105

Updated information and services can be found at:
<http://ebm.bmj.com/content/7/4/105>

	<i>These include:</i>
References	This article cites 4 articles, 1 of which you can access for free at: http://ebm.bmj.com/content/7/4/105#BIBL
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiologic studies](#) (1092)
[Pain \(anaesthesia\)](#) (75)
[Pregnancy](#) (333)
[Pain \(neurology\)](#) (413)
[Pain \(palliative care\)](#) (98)
[Childhood nutrition](#) (101)
[Clinical trials \(epidemiology\)](#) (1595)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>