


Evidence-Based Medicine



Contents

June 2008 Vol 13 No 3

Purpose and procedure

65

EBM notebook

- 66 Practice corner: managing trichotillomania—no need to pull your hair out
- 67 Interpreting meta-analysis in systematic reviews

Letter

- 69 Proactive monitoring for newly emerging evidence: The lost step of EBP?

Evidently

70

Therapeutics

- 71 Testosterone supplementation did not prevent cognitive decline or increase bone mineral density in older men
- 72  Review: use of pedometers increases physical activity in adults
- 73 Review: combination therapy with renin angiotensin inhibitors reduces proteinuria more than single drugs alone in renal disease
- 74 Review: corticosteroids plus LABAs reduce exacerbations and mortality more than steroids alone in COPD
- 75 Adjunctive diclofenac and spinal manipulation did not speed recovery of acute low back pain
- 76 Review: individual DMARDs have similar efficacy for RA, but combination therapy improves response
- 77 Adjunctive prandial or biphasic insulin for type 2 diabetes improved glycaemic control but increased hypoglycaemia and weight gain
- 78 A patient-adjusted insulin algorithm was non-inferior to standard of care for glycaemic control in type 2 diabetes
- 79 Review: long-acting insulin analogues do not improve glycaemic control but reduce nocturnal hypoglycaemia
- 80 5 days of nitrofurantoin was equivalent to 3 days of trimethoprim-sulfamethoxazole for women with non-complicated cystitis

- 81 Review: multiple doses of antenatal corticosteroids prevent preterm neonatal respiratory disease more than a single dose
- 82 Review: misoprostol and intramuscular prostaglandins do not prevent postpartum haemorrhage more than injectable uterotonics
- 83 Review: calcium supplementation during pregnancy reduces the risk of pre-eclampsia

Diagnosis

- 84 Random urine protein:creatinine ratio was an accurate method for diagnosing proteinuria in pregnant women with hypertension
- 85 Review: non-psychiatric physicians have low accuracy for recognising depression in their patients

Prognosis

- 86 Review: risk of stroke is high at 2, 30, and 90 days after transient ischaemic attack

Aetiology

- 87 Review: lower total cholesterol is associated with reduced risk of death from ischaemic heart disease but not stroke
- 88 Review: silent brain infarcts are common in the elderly general population
- 89 Glomerular filtration rate < 45 ml/min/1.73 m² with microalbuminuria was associated with increased CV death
- 90 BMI and cardiorespiratory fitness predicted mortality in older adults

Resource reviews

92

Other articles noted

95

Glossary

Inside back cover

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.



www.publicationethics.org.uk
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics