

Evidence-Based Medicine



Contents

Purpose and procedure

1

Editorial

- 2 New look Evidence-Based Medicine



Therapeutics

- 3 Simple physical interventions such as hand washing and wearing masks can reduce spread of respiratory viruses
- 4 Combined educational and contraceptive interventions reduce unplanned teenage pregnancy, but how useful is this finding for policy and practice?
- 5 Feedback to internal medicine ward physicians from multidisciplinary antimicrobial use teams improves appropriate antimicrobial use
- 6 No difference between short-course and long-course antibiotics for bacterial meningitis in children, but available evidence limited
- 7 High efficacy of HPV 16/18 AS04-adjuvanted vaccine shown against CIN2+ associated with HPV-16/18 and non-vaccine oncogenic HPV types
- 8 Use of inhaled budesonide for 1 year does not increase risk of pneumonia in patients with chronic obstructive pulmonary disease
- 9 "Hospital at home" care shows similar mortality and subsequent hospital admissions to hospital care for older patients with acutely decompensated chronic heart failure
- 10 Healthcare assistant case management may reduce depression symptoms in primary care patients with major depression more than usual care
- 11 Induction of labour improves maternal outcomes compared with expectant monitoring in women with gestational hypertension or mild pre-eclampsia
- 13 Immediate colposcopy referral in women with low-grade abnormal results on cervical cytology detects more CIN2 or worse lesions than cytological surveillance in care, but might lead to overtreatment
- 14 Effects of physical fitness training on mortality, independence and disability after stroke unclear

February 2010 Vol 15 No 1

- 15 Antipyretic drugs do not reduce recurrences of febrile seizures in children with previous febrile seizure
- 16 Interactive booklet reduces antibiotic prescribing for respiratory tract infections in children, but not parent satisfaction
- 17 Web-based screening and brief motivational intervention reduces alcohol use in heavy-drinking undergraduates at up to 6 months
- 18 Preventive antibiotics reduce infections but not mortality in adults with acute stroke
- 19 Adding atypical antipsychotics to antidepressants increases response in treatment-resistant major depression but increases discontinuation as a result of adverse events
- 20 Neuraminidase inhibitors produce a small reduction in duration of seasonal influenza in children and reduce transmission in affected households, but effects on serious complications unclear
- 21 Knee strengthening exercises at home, but not dietary intervention, reduce knee pain and improve knee function in overweight and obese people over 2 years
- 23 Primary care screening and brief counselling for overweight or mildly obese children does not improve BMI, nutrition or physical activity
- 24 Corticosteroids added to antibiotic therapy aid pain resolution in adults with sore throat, particularly severe or exudative sore throat
- 25 Cochrane Review highlights the need for more targeted research on the tolerability of malaria chemoprophylaxis in travellers
- 26 No difference in peak troponin level in immediate versus delayed (next working day) intervention for acute coronary syndrome without ST-segment elevation

Diagnosis

- 28 Point-of-care D-dimer tests can contribute to patient management in outpatients with suspected venous thromboembolism, particularly those at low risk
- 29 GPs good at ruling out depression in primary care, but false positives common

Reprinted from DTB

- 31 Aspirin for primary prevention of cardiovascular disease?



This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator
network



When you have finished
with this please recycle it