

Evidence-Based Medicine



Contents

Editorial

- 169** Why we need more breast cancer screening trials



Systematic review

- 171** Thiazolidinediones for plaque psoriasis: a systematic review and meta-analysis

Therapeutics

- 177** Acute conjunctivitis in primary care: antibiotics and placebo associated with small increase in the proportion cured by 7 days compared with no treatment
- 178** In patients with coronary artery disease and heart failure, the addition of coronary-artery bypass grafting to medical therapy has no effect on the overall risk of death from any cause, but assessment of suitability for surgery may predict survival benefit in subgroups
- 179** Early umbilical cord clamping increases the risk of neonatal anaemia and infant iron deficiency
- 181** For people with persistent tension-type neck pain, a multicomponent pain and stress self-management intervention gives better improvement in ability to control pain and self-efficacy, but not disability, than physical therapy
- 182** Catheter-directed thrombolysis prevents post-thrombotic syndrome in patients with acute deep vein thrombosis in the upper half of the thigh
- 183** The legacy effect: 4.5 years of a chlorthalidone-based antihypertensive regimen reduces cardiovascular mortality and prolongs cardiovascular disease-free survival at 22 years for older patients with isolated systolic hypertension
- 185** Combined vitamin D and calcium supplementation may reduce fracture risk in institutionalised older people but has uncertain role in cancer prevention
- 186** Study of the effect of calcium supplementation during pregnancy on the risk of preterm birth or low birthweight infants is inconclusive
- 187** Nicotine replacement therapy increases quit attempts among unmotivated smokers when added to encouragement to practice quitting
- 188** Forty-month trial suggests repeated hyaluronic acid injections for people with knee osteoarthritis may act as a long-term slow acting drug

Prognosis

- 190** Of 3700 children thought to have non-cardiac chest pain at initial paediatric cardiology clinic evaluation, none suffered cardiac death over a median of 4 years follow-up

December 2012 Vol 17 No 6

Aetiology

- 192** Statin use in postmenopausal women is associated with an increased risk of incident diabetes mellitus
- 193** Night shift work increases the risk for type 2 diabetes

Economic analysis

- 195** Screening for postmenopausal osteoporosis with any modality is cost-effective including screening initiation at 55 years of age

PostScript

- 197** Response to Drs Hoberman and Paradise
- 198** Is recent advice on exclusive breastfeeding consistent with the data presented?
- 198** Correction

Electronic pages

- e6** Purpose and procedure
- e7** Higher breastfeeding intensity associated with improved postpartum glucose metabolism in women with recent gestational diabetes
- e8** For women with term low-risk pregnancies, risk of perinatal mortality and intrapartum morbidity similar across planned places of birth
- e9** In China, hospital births are associated with a 62% decreased risk of neonatal mortality
- e10** Serial changes in highly sensitive cardiac troponin improve the early diagnosis of acute myocardial infarction
- e11** Children with minor blunt head trauma, a Glasgow Coma Scale score of 14 or 15 and a normal CT scan are at very low risk of traumatic findings on subsequent neuroimaging or of requiring neurosurgical intervention
- e12** Low-molecular-weight heparin prophylaxis does not affect mortality in acutely ill medical patients at low risk for venous thromboembolism
- e13** Multicondition collaborative care intervention for people with coronary heart disease and/or diabetes, depression and poor control of hypertension, blood sugar or hypercholesterolemia improves disability and quality of life compared with usual care
- e14** Self-monitoring of oral anticoagulation reduces thromboembolic events and does not increase risk of bleeding in selected patients and settings
- e15** Topical dexamethasone for recurrent aphthous ulceration reduces pain and size and increases healing with no significant adverse events
- e16** Antibiotics provide no clinically important benefit in mild to moderate acute sinusitis



This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator network



When you have finished with this please recycle it