

Evidence-Based Medicine



Contents

April 2016 Volume 21 No 2

Perspective

- 41 **'Cognitive biases plus': covert subverters of healthcare evidence**



Original EBM research

- 46 Evidence-based medicine and life-long learning competency requirements in new residency teaching standards
- 47 **Ophthalmologist knowledge of evidence-based medicine and clinical practice guideline recommendations**

Methods

- 55 Importance and methods of searching for E-publications ahead of print in systematic reviews

Therapeutics/Prevention

- 60 Adjunctive corticosteroids improve the need for mechanical ventilation and shorten hospital duration in patients hospitalised with community-acquired pneumonia
- 61 A modified Valsalva manoeuvre results in greater termination of supraventricular tachycardia than standard Valsalva manoeuvre
- 62 Ibuprofen for the treatment of TTH
- 63 Best evidence supports topical corticosteroids and routine saline irrigations for medical treatment of chronic rhinosinusitis
- 64 Artemisinin-based combination therapy (ACT) versus atovaquone-proguanil: do not choose between but, rather, combine them
- 65 Autoinflation reduces middle ear effusion in children with otitis media with effusion
- 66 **Low-molecular-weight heparin for the treatment of acute venous thromboembolism in patients with active cancer**
- 67 Non-toxicogenic *Clostridium difficile* to prevent recurrent *C. difficile* infection

- 68 Prescribing of metformin based on estimated GFR rather than serum creatinine expands the eligible population and is likely safe
- 69 Four sessions of spinal manipulation, simple exercises and education are not better than usual care for patients with acute low back pain
- 70 Aerobic exercise training reduces bronchial hyper-responsiveness and serum pro-inflammatory cytokines in patients with asthma
- 71 Ultrasound-guided corticosteroid injections for adhesive capsulitis more effective than placebo
- 72 Mediterranean diet supplemented with extra virgin olive oil reduces the incidence of invasive breast cancer in a randomised controlled trial
- 73 Complete revascularisation in patients with ST-segment elevation myocardial infarction and multivessel disease: contemporary data in context
- 75 Maternal postpartum high-dose vitamin D3 supplementation (6400 IU/day) or conventional infant vitamin D3 supplementation (400 IU/day) lead to similar vitamin D status of healthy exclusively/fully breastfeeding infants by 7 months of age

Prognosis

- 76 Increasing children's time spent outdoors reduces the incidence of myopia

Aetiology/Harm

- 77 Light to moderate alcohol intake may be associated with non-minimal cancer risk
- 78 More data needed on hysteroscopic compared with laparoscopic sterilisation
- 79 Emergency department visits and hospitalisations for adverse events related to dietary supplements are common
- 80 Early exposure to dogs and farm animals reduces risk of childhood asthma

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
Member since 2010
JM0659
www.publicationethics.org.uk

equator network

recycle
When you have finished with this please recycle it

Receive regular table of contents by email.
Register using this QR code.

