


# Evidence-Based Medicine



## Contents

### EBM opinion and debate

- 157** Meta-analysis using individual participant data from randomised trials: opportunities and limitations created by access to raw data

### Maternal and child health

- 162** USPSTF is unable to recommend for or against lipid screening to identify multifactorial dyslipidaemia in childhood: no recommendation is not without consequences

### Original EBM research

- 164** What does expert opinion in guidelines mean? a meta-epidemiological study

### Evidence synthesis

#### General medicine

- 170** Quality of reporting of harms in randomised controlled trials of pharmacological interventions for rheumatoid arthritis: a systematic review

### Commentaries

#### Primary care

- 178** Longer screening intervals are recommended following a negative HPV test in primary cervical screening
- 179** Haem iron and nitrate/nitrite account for much of the mortality increase associated with red meat consumption
- 180** Value of screening for and treating pre-diabetes is reduced by low test accuracy and indirect evidence of impact on patient important outcomes

#### General medicine

- 182** Caffeine does not increase the arrhythmic burden in patients with heart failure and left ventricular systolic dysfunction
- 183** In stable COPD, long-acting muscarinic antagonist plus long-acting beta-agonists resulted in less exacerbations, pneumonia and larger improvement in FEV<sub>1</sub> than long-acting beta-agonists plus inhaled corticosteroids

## October 2017 Volume 22 Issue 5

- 184** CETP inhibition improves the lipid profile but has no effect on clinical cardiovascular outcomes in high-risk patients

- 185** Early renin-angiotensin system inhibition induced renal deterioration may be a predictor for long-term cardiorenal outcomes

### Maternal and child health

- 187** Antibiotic utilisation in very low birth weight infants without sepsis or necrotising enterocolitis is associated with multiple adverse outcomes
- 188** Serial transvaginal cervical length measurements and quantitative vaginal fetal fibronectin concentrations did not predict spontaneous preterm birth in low-risk nulliparous women
- 189** Antenatal corticosteroid administration between 24 hours and 7 days before extremely preterm delivery is associated with the lowest rate of mortality

### Emergency care

- 190** Routine invasive strategies compared with conservative strategies do not lower the all-cause mortality in patients with non-ST elevation myocardial infarction and unstable angina

### Surgery


- 191** Over half of the patients who undergo adjustable gastric banding may require revision bariatric surgery
- 192** In localised prostate cancer, radical prostatectomy was associated with more sexual dysfunction and urinary incontinence than radiation or active surveillance
- 193** Meniscal resection may not benefit patients with traumatic meniscal tears


### Mental health

- 194** Parent-delivered CBT may reduce intervention cost, but questions arise about effectiveness

### Letters

- 195** Effectiveness of interventions based on implantable devices: meta-analyses or systematic reviews that fail to indicate which device brands were used
- 196** Comparing levels of evidence between Choosing Wisely and Essential Evidence Plus

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

 This journal is a member of and subscribes to the principles of the Committee on Publication Ethics [www.publicationethics.org.uk](http://www.publicationethics.org.uk)

Member since 2010  
JMO6659

 equator network

 recycle  
When you have finished with this please recycle it