

Is there a smoker's paradox in COVID-19?

Muhammad Shariq Usman ¹, Tariq Jamal Siddiqi,¹
Muhammad Shahzeb Khan,² Urvish K Patel,³ Izza Shahid,⁴
Jawad Ahmed,¹ Ankur Kalra,⁵ Erin D Michos⁶

10.1136/bmjebm-2020-111492

¹Department of Internal Medicine, Dow University of Health Sciences, Karachi, Pakistan

²Department of Internal Medicine, John H Stroger Hospital of Cook County, Chicago, Illinois, USA

³Department of Public Health and Neurology, Icahn School of Medicine at Mount Sinai, New York City, New York, USA

⁴Department of Internal Medicine, Ziauddin University, Karachi, Pakistan

⁵Department of Cardiovascular Medicine, Cleveland Clinic, Cleveland, Ohio, USA

⁶Department of Cardiovascular Diseases, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

Correspondence to:

Dr Muhammad Shariq Usman, Internal Medicine, Dow University of Health Sciences, Karachi, SD 74200, Pakistan; shariqusman@outlook.com



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Usman MS, Siddiqi TJ, Khan MS, et al. *BMJ Evidence-Based Medicine* 2021;26:279–284.

Abstract

Although it is well established that cigarette smoking is associated with morbidity and mortality in several respiratory infections, data from recent studies suggest that active smokers are underrepresented among patients with COVID-19. This has led to claims that a 'smoker's paradox' may exist in COVID-19, wherein smokers are protected from infection and severe complications of COVID-19. We aimed to review and summarise existing literature in this context. Electronic databases were searched for articles that reported prevalence of smokers among patients with COVID-19 or studied any association of smoking with outcomes among patients with COVID-19. We identified several biases and knowledge gaps which may give the false impression that smoking is protective in COVID-19. As of now, the data supporting smoker's paradox claims are limited and questionable. Plausible biologic mechanisms by which smoking might be protective in COVID-19 include an anti-inflammatory effect of nicotine, a blunted immune response in smokers (reducing the risk of a cytokine storm in COVID-19) and increased nitric oxide in the respiratory tract (which may inhibit replication of SARS-CoV-2 and its entry into cells). On the other hand, smoking may worsen susceptibility and prognosis in COVID-19, in a manner similar to other respiratory infections. The claims of a protective effect must be viewed with extreme caution by both the general population as well as clinicians. Further investigations into the interaction between smoking and COVID-19 are warranted to accurately assess the risk of contracting COVID-19 among smokers, and progression to mechanical ventilation or death in patients suffering from it.

Background

The relationship between smoking and coronavirus disease 2019 (COVID-19) has recently received a lot of attention. It is well established that cigarette smoking predisposes to several respiratory infections and is also associated with relatively poor outcomes in those suffering from these infections.¹ Based on this knowledge, the World Health Organization (WHO) and the U.S. Food and Drug Administration (FDA) both released statements warning that smoking may increase the risk and severity of COVID-19.^{2,3} The assertion of risk associated with smoking, however, was rooted in expectation rather than data. Interestingly, recent studies have

shown that active smokers are actually underrepresented among those requiring hospital treatment for COVID-19 in several countries including the USA, China and France.⁴ This led to widespread claims that smoking may be protective against COVID-19.^{4,5} Although a protective effect of smoking in COVID-19 may seem counterintuitive, researchers have suggested several mechanisms by which this 'smoker's paradox' may be possible.⁴ In this article, we aim to (1) review currently available data regarding the relationship between smoking and COVID-19, (2) highlight possible biases in the available data and discuss current knowledge gaps, and (3) summarise proposed mechanisms of interaction between smoking and COVID-19.

Smoker's paradox

The concept of 'smoker's paradox' is not new. The term was first coined in 1995 to describe the unexpected favourable outcome of decreased short-term mortality in smokers after acute coronary syndrome^{6,7} and stroke,⁸ when compared with non-smokers. Conflicting theories exist in the literature, where some have proposed that this finding may be due to a decreased inflammatory response in smokers. Others suggest that this may be attributed to the difference in pathogenesis of myocardial infarction (MI) between the two groups, where a thrombotic MI occurring in smokers, compared with an atherosclerotic plaque-mediated event in non-smokers, would respond better to thrombolytic interventions.⁷ Another thought was that smokers may have ischaemic pre-conditioning, which would reduce infarct size. On the other hand, many simply attributed the apparently lower short-term mortality in smokers to differences in baseline characteristics between the two groups (such as younger age and fewer cardiovascular risk factors at the time of MI in smokers)⁹ or residual confounding by other factors (such as age at initiation of smoking)¹⁰ which are often not adequately adjusted for in observational studies.¹¹ Of note, the majority of studies reporting this phenomenon group both current and former smokers under the common classification of 'smoking', despite smoking cessation showing a significant decrease in cardiovascular disease and subsequent mortality after a couple of years.¹² Grouping both current and former smokers together can have potential confounding effects of nicotine withdrawal on the results, while excluding former smokers leads to ambiguity surrounding the existence of a smoker's paradox in this subgroup.

A recent study on acute ST-elevation MI patients failed to confirm this hypothesised benefit of smoking, and instead showed worse outcomes in smokers after MI.¹³ Hence, the idea of a smoker's paradox in cardiovascular disease has lost momentum in recent years. Given the recent claims that smoking has a protective effect in COVID-19, we must entertain the possibility of a smoker's paradox in COVID-19. However, the experience with cardiovascular disease must be kept in mind, as many of the same biases may apply in the current situation.

Current evidence

The causal effects of smoking on disease processes have traditionally been difficult to study due to the impracticality of randomised controlled trials and a wide range of potential confounding factors. In such cases, causality is usually inferred via the process of triangulation, which involves using several research methods to study the same phenomenon. Thus, the same must be applied to study the relationship between smoking and COVID-19.

Search strategy and selection criteria of the reviewed studies

The PubMed and Scopus databases were searched in May 2020. Different combinations of following keywords and their MeSH terms were used: 'Coronavirus disease 2019', 'COVID-19', 'characteristics', 'smoking', 'nicotine' and 'tobacco'. Bibliographies of included articles were also hand-searched for any articles that may have potentially been missed. In addition, a cursory search of pre-print servers was also performed for relevant articles with prominent results. Studies were included in this review if they reported smoking prevalence and/or its association with clinical outcomes in patients with diagnosed COVID-19. Studies reporting original data and systematic reviews/meta-analyses were considered. Table 1 summarises the characteristics of the studies reviewed for the purpose of this article. Furthermore, web searches (on www.google.co.uk) were conducted for relevant news articles and media reports.

Prevalence of smokers among the COVID-19 population

The low prevalence of active smokers among hospitalised patients with COVID-19 has been a consistent finding across most published studies.^{14–21} A recent meta-analysis of 13 studies (n=5960 patients) demonstrated that the pooled prevalence of smokers among hospitalised patients with COVID-19 in China was 6.5% (95% CI 4.9% to 8.2%).²² This number is drastically lower than the prevalence of active smokers in the general population of China (26.6%).²³ A recent study from France (currently available on a pre-print server) demonstrated a lower incidence of active smokers among 482 symptomatic patients with COVID-19, when compared with the general population.²⁴ This French study included both inpatients and outpatients; thus, their results suggest that smokers may be less likely to be infected with COVID-19. Evidence from three published studies from the USA, all reporting data on hospitalised patients with COVID-19 in New York City, have reported different prevalence of active smokers. The prevalence of active smokers in the US population is 13.7%²⁵; however, in the study by Goyal *et al*,²⁶ smokers made up 20/393 (5.1%) of the hospitalised population. Cummings *et al*²⁷ reported an aggregated prevalence of current and former smokers in critically ill patients (33/257; 12.8%), which was also relatively low. The study by Richardson *et al* demonstrated a considerably higher prevalence (558/3567; 15.6%).²⁸ To the best of our knowledge, the study by Richardson

and colleagues is the only one to date in which the prevalence of smokers resembles that of the general population.

Association of smoking with clinical outcomes in COVID-19

Although smokers appear to be consistently underrepresented among patients with COVID-19, the link between smoking and clinical outcomes in these patients remains less clear. A systematic review, published in March 2020 by Vardavas and Nikitara,²⁹ concluded that smoking was associated with disease progression and increased adverse outcomes in COVID-19; however, this conclusion was based on limited and mostly unadjusted data. In a meta-analysis published in the same time period, Lippi and Henry³⁰ demonstrated no significant association between active smoking and COVID-19 severity (OR 1.69 (95% CI 0.41 to 6.92); p=0.254). The wide CIs derived in the meta-analysis are an indication of the heterogeneity of the studies and low reliability of the data. A large study of the general population of the UK found that smoking was significantly associated with increased COVID-19 mortality after age and sex adjustment (OR 1.25 (1.12 to 1.40)).³¹ However, after adjustment for multiple additional covariates, the same study found that smoking was associated with a reduced risk for COVID-19 mortality (OR 0.88 (0.79 to 0.99)). This protective association remained consistent after several individual adjustments to the model. Clearly, current evidence in the area is equivocal and limited, and further investigation is needed.

Limitations of current evidence

Several possible biases must be taken into consideration when assessing the current evidence in the area. First, several concerns have been raised regarding the quality of the data reported in published COVID-19 papers. Early in the pandemic crisis, there was a race to publish, which still continues to some extent. This likely resulted in aberrant and non-standardised data collection methods, poor statistical analysis, and limited scrutiny during editorial and peer review. In the context of smoking and COVID-19, poor data collection can lead to several erroneous conclusions. If patients with missing smoking data are not eliminated from the total pool, smokers may be wrongly underrepresented. Furthermore, it is difficult to get accurate history from patients who are either intubated or in respiratory failure. If data from these patients are missing, and these patients are not removed from the denominator, it can give a false impression that smokers are less likely to develop severe disease. Such errors are not far-fetched, especially when evaluating the effect of smoking is not the primary goal of the study. Second, it must be noted that most published studies have not reported the duration (years) or frequency (number of cigarettes) of smoking, hence these cannot be accounted for. Furthermore, most studies did not report the number of former smokers. It is conceivable that a proportion of the patients quit smoking only in the recent past (eg, after the start of the pandemic or after being diagnosed with COVID-19). As discussed earlier, inclusion of both former and current smokers as 'smokers' in these studies can create biases that contribute to smoker's paradox. Third, none of the studies used objective biomarkers to determine active smoking. In fact, the source of smoking history in most published studies remains almost totally uncertain. Lastly, it must be kept in mind that the prevalence data in most studies are unadjusted. While using the general population as a control is convenient and offers some insight, the COVID-19 population likely has a different distribution of age, sex, comorbidities and smoking patterns.

Table 1 Characteristics of studies reviewed

Study	Country	Sample size, N	Diabetes, N (%)	Hypertension, N (%)	CVD, N (%)	Pulmonary disease, N (%)	Active smokers, N (%)	Former smokers, N (%)	ICU admission, N (%)	Mortality, N (%)	Notes
Original studies											
Huang <i>et al</i> ¹⁴	China	41	8 (19.5)	6 (14.6)	6 (14.6)	1 (2.4)	3 (7.3)	NR	7 (17.1)	6 (14.6)	–
Guan <i>et al</i> ¹⁵	China	1085	81 (7.5)	165 (15.2)	27 (2.5)	12 (1.1)	137 (12.6)	21 (1.9)	55 (5.0)	15 (1.4)	–
Zhang <i>et al</i> ¹⁶	China	140	17 (12.1)	42 (30.0)	12 (8.6)	2 (1.4)	2 (1.4)	7 (5.0)	–	–	–
Yang <i>et al</i> ¹⁷	China	52	9 (17.3)	–	5 (9.6)	4 (7.7)	2 (3.8)	NR	52 (100.0)	32 (61.5)	–
Mo <i>et al</i> ¹⁸	China	155	15 (9.7)	37 (21.9)	15 (9.7)	5 (3.2)	6 (3.9)	NR	–	–	–
Zhou <i>et al</i> ¹⁹	China	191	36 (18.9)	58 (30.4)	15 (7.9)	6 (3.1)	11 (5.8)	NR	50 (26.2)	54 (28.3)	–
Guo <i>et al</i> ²⁰	China	187	28 (14.9)	61 (32.6)	21 (11.2)	–	18 (9.6)	NR	–	43 (23.0)	–
Richardson <i>et al</i> ²⁸	USA	3567	1808 (50.7)	3026 (84.8)	966 (27.1)	920 (25.8)	558 (15.6)	NR	1281 (22.5)	553 (9.7)	–
Goyal <i>et al</i> ²⁶	USA	393	99 (25.2)	197 (50.1)	82 (20.9)	69 (17.6)	20 (5.1)	NR	–	40 (10.2)	–
Qin <i>et al</i> ²¹	China	452	75 (16.6)	135 (29.9)	27 (5.9)	12 (2.7)	7 (1.5)	NR	–	–	–
Cummings <i>et al</i> ²⁷	USA	257	92 (35.8)	162 (63.0)	49 (19.1)	24 (9.3)	33 (12.8)*	NR	203 (79.0)	101 (39.3)	–
Meta-analyses											
Farsalinos <i>et al</i> ²²	China	5960	–	–	–	–	450 (7.6)	35 (0.6)	–	–	Pooled prevalence of smokers among hospitalised patients with COVID-19 in China was 6.5% (95% CI 4.9% to 8.2%)
Lippi <i>et al</i> ⁴⁰	China	1399	–	–	–	–	–	–	–	–	No significant association between active smoking and COVID-19 severity (OR 1.69 (0.41 to 6.92); p=0.254)
Preprints											
Miyara <i>et al</i> ²⁴	France	482	102/473	158/473	–	29/473	34/473	285/473	–	–	Daily smokers have lower probability of developing severe COVID-19 when compared with the general population
OpenSAFELY Collaborators ³¹	UK	17 425 455	489 297 (2.8)	626 122 (34.2)	1173 433 (6.7)	707 284 (4.1)	962 373 (17.0)	737 545 (32.9)	–	5683 (0.03)	On adjustment for multiple covariates, smoking was associated with a reduced risk for COVID-19 mortality (OR 0.88 (0.79 to 0.99))

*Study reported 'current or former smokers'.

CI, Confidence interval; CVD, cardiovascular disease; ICU, intensive care unit; NR, not reported; OR, Odds ratio; UK, United Kingdom; USA, United States of America.

Proposed interactions between smoking and COVID-19

Mechanisms by which smoking may have a protective effect

Despite all the aforementioned concerns with residual confounding, there are a few plausible biologic mechanisms by which smoking might indeed be associated with less severe COVID-19 disease. First, nicotine, which is an agonist of the cholinergic anti-inflammatory pathway, could potentially offer a protective effect in COVID-19.³² This is also a potential explanation for its ability to treat inflamed bowels. It inhibits production of pro-inflammatory cytokines such as TNF, IL-1 and IL-6, without inhibiting the anti-inflammatory cytokines such as IL-10,^{32 33} thereby possibly protecting against the cytokine-storm syndrome, a phenomenon responsible for the pathophysiology of severe COVID-19.³⁴ Continuous suppression of these systemic cytokines in smokers may adapt their immune response in becoming more tolerant and therefore less reactive to the continuous inflammatory stimuli when compared with patients who have never smoked.³⁵ Second, nitric oxide (NO) produced during smoking is involved in maintaining airway dilation and filtration prior to its entry in lungs.³⁶ NO has been shown to inhibit the replication of SARS-CoV-2 and

its entry into cells.³⁷ Third, smoking may upregulate ACE2, an anti-inflammatory protein, in lower respiratory tract—as further discussed below (figure 1).

Mechanisms by which smoking may have a harmful effect

Smokers are almost certainly more vulnerable to initial exposure to the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus responsible for COVID-19, due to repetitive hand-to-mouth movements and contact of possibly contaminated smoking apparatus with lips.³⁸ Moreover, cigarettes are also sometimes shared between individuals. Patients may initially be asymptomatic, and such sharing may lead to inadvertent dissemination of the virus. Furthermore, smoking is a strong risk factor for premature cardiovascular diseases and chronic obstructive pulmonary disease, which have been shown to be associated with increased COVID-19 severity and mortality.^{39 40} Tobacco smoke suppresses the function of innate immune cells, including respiratory epithelium, alveolar surfactant, macrophage, neutrophils and lymphocytes. Tobacco contains components which disrupt the normal epithelial lining of the respiratory system leading to




Figure 1 Proposed mechanisms of interaction between smoking and COVID-19. COPD, chronic obstructive pulmonary disease; ACE2, Angiotensin converting enzyme 2; IL-1, interleukin 1; IL-6, interleukin 6; NO, nitric oxide; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; TNF, tumor necrosis factor

increased oxidative injury and impairment of mucociliary clearance.⁴¹ This may make smokers more susceptible to developing complications of COVID-19, such as pneumonia. Smoking also impairs production of surfactant, thereby leading to decreased leucocyte function and impaired host immunity.⁴² Moreover, the phagocytic function of alveolar macrophage is also significantly reduced by smoking, causing decreased clearance of inflammatory cells and debris from the lungs.⁴³ Furthermore, cigarette smoking can also alter T-cell responses, which can lead to increased susceptibility to respiratory tract infections, which can be particularly detrimental in patients with concomitant COVID-19.⁴³

Smoking and ACE2

ACE2, a cell surface protein expressed on several human tissues, is a known receptor for the SARS-CoV-2 virus. The physiological function of ACE2 is degradation of angiotensin II to angiotensin₁₋₇, which serves as a protective function against overactivity of angiotensin II.⁴⁴ In different experimental models, angiotensin II overactivity is reported to have triggered myocardial dysfunction, interstitial fibrosis, endothelial dysfunction, inflammation, oxidative stress and coagulation in addition to its primary function as a potent vasoconstrictor.⁴⁵ On entering cells via ACE2, the virus then downregulates this protein.⁴⁴ This prevents degradation of angiotensin II, and the subsequent local angiotensin II overactivity may be responsible for many of the inflammatory and thrombotic complications of COVID-19.⁴⁴

Although it has been proposed that smoking modulates the expression of ACE2, the precise effect remains unclear. Recent literature reports that smoking and nicotine may upregulate ACE2,⁴⁶ while studies published prior to the COVID-19 pandemic report smoking and nicotine to contribute to downregulation of ACE2.⁴⁷ If smoking does in fact upregulate ACE2, then it may serve as a protective factor in decreasing disease severity. However if smoking diminishes ACE2 expression, further downregulation by SARS-CoV-2 virus may amplify the disease severity and promote adverse outcomes in this cohort.⁴⁴ On the contrary, increased ACE2 receptors in smokers also potentially leads to increased viral receptors through which SARS-CoV-2 virus may enter the body.³⁸ This contradiction warrants more research in order to most accurately determine the effect of nicotine on ACE2 expression and downstream consequences.

Conclusion

Although, on the surface, it appears that smokers are underrepresented among the COVID-19 population, we highlight several biases and knowledge gaps which have yet to be addressed. Currently reported data are questionable, and a protective effect should not be inferred. Claims that smoking is protective in COVID-19 can be detrimental to public health and should be viewed with extreme caution by both the general population as well as clinicians. Even if smoking did offer a protective effect in COVID-19, this is unlikely to outweigh the numerous proven adverse health effects of smoking. Hypotheses for both protective and detrimental effects of smoking on COVID-19 have been proposed in the literature (figure 1). Further investigation into the interaction between smoking and COVID-19 is warranted for two main reasons: (1) to assess the possibility of nicotine as a therapeutic option; (2) to allow accurate assessment of risk of contracting COVID-19 among smokers, and progression to mechanical ventilation or death in patients suffering from it.

Contributors MSU, TJS, AK and EDM contributed to conception of study and MSK and UKP contributed to study design. IS

and JA contributed to literature search and MSU, TJS and MSK contributed to study selection. MSU, TJS, IS and JA contributed to initial manuscript draft. MSK, UKP, AK and EDM critically revised the manuscript for important intellectual content. MSU and TJS have contributed equally and are co-first authors. All authors have read and approved the final version of manuscript and agree to be accountable for accuracy and integrity of all aspects of the work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval No human participants were involved in the study and it uses publicly available data.

Provenance and peer review Not commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iD

Muhammad Shariq Usman <http://orcid.org/0000-0001-9747-8892>

References

- 1 Arcavi L, Benowitz NL. Cigarette smoking and infection. *Arch Intern Med* 2004;164:2206–16.
- 2 World Health Organisation. Smoking and COVID-19. Available: https://www.who.int/news-room/commentaries/detail/smoking-and-covid-19?fbclid=IwAR0NmuZsrnF6kMUKM-0KMes4KPmA0sGqtsZQFi5osWC8fxeUCn_NfTo4j8U [Accessed 28 May 2020].
- 3 U.S Food and Drug Administration. Coronavirus (COVID-19) update: daily roundup, 2020. Available: https://www.fda.gov/news-events/press-announcements/coronavirus-covid-19-update-daily-roundup-april-28-2020?fbclid=IwAR0qhdtI0YuVmt7a2-nXKDwMWkzKYSbWMFQFhtFfm057er_rINN8zG3j5o [Accessed 28 May 2020].
- 4 The Economist. Smokers seem less likely than non-smokers to fall ill with covid-19. Available: <https://www.economist.com/science-and-technology/2020/05/02/smokers-seem-less-likely-than-non-smokers-to-fall-ill-with-covid-19> [Accessed 20 May 2020].
- 5 Hospital Healthcare Europe. Are smokers protected from COVID-19? Available: https://hospitalhealthcare.com/covid-19/are-smokers-protected-from-covid-19/?fbclid=IwAR0JmsHRIH8Sr-XDqAsrbhzb6PpBEYm_bnG0cewqw29cy-BaFieVKsXRdKA [Accessed Last accessed 4 Jun 2020].
- 6 Aune E, Røislien J, Mathisen M, *et al.* The “smoker's paradox” in patients with acute coronary syndrome: a systematic review. *BMC Med* 2011;9:97.
- 7 Grines CL, Topol EJ, O'Neill WW, *et al.* Effect of cigarette smoking on outcome after thrombolytic therapy for myocardial infarction. *Circulation* 1995;91:298–303.
- 8 Hussein HM, Niemann N, Parker ED, *et al.* Searching for the smoker's paradox in acute stroke patients treated with intravenous thrombolysis. *Nicotine Tob Res* 2017;19:871–6.
- 9 Fonarow GC, Abraham WT, Albert NM, *et al.* A smoker's paradox in patients hospitalized for heart failure: findings from OPTIMIZE-HF. *Eur Heart J* 2008;29:1983–91.
- 10 Fa-Binefa M, Clará A, Pérez-Fernández S, *et al.* Early smoking-onset age and risk of cardiovascular disease and mortality. *Prev Med* 2019;124:17–22.

- 11 Fewell Z, Davey Smith G, Sterne JAC. The impact of residual and unmeasured confounding in epidemiologic studies: a simulation study. *Am J Epidemiol* 2007;166:646–55.
- 12 Bakhru A, Erlinger TP. Smoking cessation and cardiovascular disease risk factors: results from the Third National Health and Nutrition Examination Survey. *PLoS Med* 2005;2:e160.
- 13 Redfors B, Furer A, Selker HP, *et al.* Effect of smoking on outcomes of primary PCI in patients with STEMI. *J Am Coll Cardiol* 2020;75:1743–54.
- 14 Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.
- 15 Guan W-J, Ni Z-Y, Hu Y, *et al.* Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382:1708–20.
- 16 Zhang J-J, Dong X, Cao Y-Y, *et al.* Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 2020;75:1730–41.
- 17 Yang X, Yu Y, Xu J, *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020;8:475–81.
- 18 Mo P, Xing Y, Xiao Y, *et al.* Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa270. [Epub ahead of print: 16 Mar 2020].
- 19 Zhou F, Yu T, Du R, *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–62.
- 20 Guo T, Fan Y, Chen M, *et al.* Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020;5:811.
- 21 Qin C, Zhou L, Hu Z, *et al.* Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis* 2020;71:762–8.
- 22 Farsalinos K, Barbouni A, Niaura R. Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: could nicotine be a therapeutic option? *Intern Emerg Med* 2020:1–8.
- 23 World Health Organization. Global Adult Tobacco Survey (GATS). Fact sheet China, 2018. Available: https://www.who.int/docs/default-source/wpro-documents/countries/china/2018-gats-china-factsheet-cn-en.pdf?sfvrsn=3f4e2da9_2 [Accessed May 2020].
- 24 Makoto Miyara FT, Valérie P, Morelot-Panzini C, *et al.* Low incidence of daily active tobacco smoking in patients with symptomatic COVID-19. *Qeios* 2020.
- 25 Centers for Disease Control and Prevention. Burden of cigarette use in the U.S. Available: <https://www.cdc.gov/tobacco/campaign/tips/resources/data/cigarette-smoking-in-united-states.html#one> [Accessed May 2020].
- 26 Goyal P, Choi JJ, Pinheiro LC. Clinical characteristics of Covid-19 in New York City. *N Engl J Med* 2020.
- 27 Cummings MJ, Baldwin MR, Abrams D, *et al.* Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* 2020.
- 28 Richardson S, Hirsch JS, Narasimhan M, *et al.* Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA* 2020;323:2052–9.
- 29 Vardavas CI, Nikitara K. COVID-19 and smoking: a systematic review of the evidence. *Tob Induc Dis* 2020;18:20.
- 30 Lippi G, Henry BM. Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19). *Eur J Intern Med* 2020;75:107–8.
- 31 Williamson EJ, Walker AJ, Bhaskaran K, *et al.* OpenSAFELY: factors associated with COVID-19 death in 17 million patients. *Nature* 2020. doi:10.1038/s41586-020-2521-4. [Epub ahead of print: 08 Jul 2020].
- 32 Wang H, Yu M, Ochani M, *et al.* Nicotinic acetylcholine receptor alpha7 subunit is an essential regulator of inflammation. *Nature* 2003;421:384–8.
- 33 Pavlov VA, Wang H, Czura CJ, *et al.* The cholinergic anti-inflammatory pathway: a missing link in neuroimmunomodulation. *Mol Med* 2003;9:125–34.
- 34 Ruan Q, Yang K, Wang W, *et al.* Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020;46:846–8.
- 35 Garufi G, Carbognin L, Orlandi A, *et al.* Smoking habit and hospitalization for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-related pneumonia: the unsolved paradox behind the evidence. *Eur J Intern Med* 2020;77:121–2.
- 36 Vleeming W, Rambali B, Opperhuizen A. The role of nitric oxide in cigarette smoking and nicotine addiction. *Nicotine Tob Res* 2002;4:341–8.
- 37 Åkerström S, Mousavi-Jazi M, Klingström J, *et al.* Nitric oxide inhibits the replication cycle of severe acute respiratory syndrome coronavirus. *J Virol* 2005;79:1966–9.
- 38 Berlin I, Thomas D, Le Faou A-L, *et al.* COVID-19 and smoking. *Nicotine Tob Res* 2020. doi:10.1093/ntr/ntaa059. [Epub ahead of print: 03 Apr 2020].
- 39 Aggarwal G, Cheruiyot I, Aggarwal S, *et al.* Association of cardiovascular disease with coronavirus disease 2019 (COVID-19) severity: a meta-analysis. *Curr Probl Cardiol* 2020;100617.
- 40 Lippi G, Henry BM. Chronic obstructive pulmonary disease is associated with severe coronavirus disease 2019 (COVID-19). *Respir Med* 2020;167:105941.
- 41 Centers for Disease Control and Prevention (US), National Center for Chronic Disease, Office on Smoking and Health (US), Prevention and Health Promotion (US). How tobacco smoke causes disease: the biology and behavioral basis for smoking-attributable disease: a report of the Surgeon General; 2010.
- 42 Honda Y, Takahashi H, Kuroki Y, *et al.* Decreased contents of surfactant proteins A and D in BAL fluids of healthy smokers. *Chest* 1996;109:1006–9.
- 43 Mehta H, Nazzal K, Sadikot RT. Cigarette smoking and innate immunity. *Inflamm Res* 2008;57:497–503.
- 44 Verdecchia P, Cavallini C, Spanevello A, *et al.* The pivotal link between ACE2 deficiency and SARS-CoV-2 infection. *Eur J Intern Med* 2020;76:14–20.
- 45 Kuba K, Imai Y, Penninger JM. Multiple functions of angiotensin-converting enzyme 2 and its relevance in cardiovascular diseases. *Circ J* 2013;77:301–8.
- 46 Brake SJ, Barnsley K, Lu W, *et al.* Smoking upregulates angiotensin-converting enzyme-2 receptor: a potential adhesion site for novel coronavirus SARS-CoV-2 (Covid-19). *J Clin Med* 2020;9. doi:10.3390/jcm9030841.
- 47 Oakes JM, Fuchs RM, Gardner JD, *et al.* Nicotine and the renin-angiotensin system. *Am J Physiol Regul Integr Comp Physiol* 2018;315:R895–906.