Evidence-Based Medicine

Contents

| Purpose and Procedu | re |
|------------------------|------------------|
| Additional Articles Al | ostracted |

EBM NOTEBOOK

| Evidence-Based Nursing: an introduction | |
|--|----|
| Using Best Evidence in clinical practice | 39 |
| Correction | 40 |

THERAPEUTICS

| Review: Lowering cholesterol levels with statin drugs reduces fatal and nonfatal strokes and total mortality 41 |
|--|
| Stroke unit care improved survival and function for 5 years after an acute stroke |
| Enoxaparin reduced combined coronary events in unstable angina and non-Q-wave MI at 14 and 30 days 43 |
| Coronary angioplasty led to more nonfatal MIs than medical therapy in patients with coronary artery disease 44 |
| Low-molecular-weight heparin and unfractionated heparin were equally effective for acute pulmonary embolism 45 |
| Indinavir plus zidovudine and lamivudine reduced HIV RNA levels in most HIV-infected persons with previous antiretroviral treatment |
| Indinavir plus zidovudine and lamivudine reduced the progression to AIDS in patients with previous zidovudine therapy |
| Antiseptic catheters reduced catheter-related infections and were well tolerated |
| A catheter coated with minocycline and rifampin reduced bloodstream infections |
| Review: Intensive insulin therapy for type 1 diabetes mellitus increases risk for severe hypoglycemia, and pump therapy increases ketoacidosis |
| Review: Noninvasive positive-pressure ventilation reduces mortality and need for intubation in the intensive care unit |

| Review: Antibiotics reduce the already low complication rate and shorten the duration of symptoms in sore throat at day 3 but not day 7 |
|---|
| Prescribing antibiotics for sore throat increased reattendance |
| Review: High-compression bandages, stockings, and Unna's boot improve healing rates in venous leg ulcers 54 |
| Review: Plasma-derived hepatitis B vaccine is safe and effective for health care workers |
| Calcium supplements during normal pregnancy did not prevent pre-eclampsia or other related disorders56 |
| Calcium and vitamin D reduced nonvertebral fractures and bone loss |

PROGNOSIS

| Bleeding occurred in one fifth of recognised pregnancies |
|--|
| before the twentieth week and over half of these |
| miscarried |

ETIOLOGY

| Childhood acute lymphoblastic leukaemia was not linked to magnetic-field exposure from power lines59 |
|---|
| Pregnancy after breast cancer treatment was not associated with an increased risk for death |
| ACE inhibitor use was associated with severe hypoglycemia in patients with diabetes mellitus61 |

QUALITY IMPROVEMENT

| Review: Provision of costing information changes physician behaviour and reduces costs | 62 |
|---|----|
| Feedback to general practitioners increased prescribing of aspirin to patients with ischaemic heart disease | 63 |

ECONOMICS

| Loss-of-life cost for paediatric illness associated with parental smoking was estimated at U.S. \$8.2 billion |
|---|
| per year |
| Glossary Inside back cover |