Evidence-Based Medicine

Contents

Purpose and Procedure 2	Alosetron was effective and safe for relieving abdominal symptoms in women with irritable bowel syndrome21
Additional articles abstracted in ACP Journal Club	Review: 3 to 7 months of prednisone is more effective than 2 months in children with steroid responsive nephrotic syndrome
EBM notebook	Diagnosis
The rocky road: qualitative research as evidence 4	Smoking, age, and laryngeal height predicted obstructive airway disease in adults23
Resource corner 7	Abdominal palpation had moderate sensitivity and specificity for detecting abdominal aortic aneurysm24
Therapeutics	Review: the accuracy of troponin T and I tests in acute myocardial infarction depends on timing after onset of symptoms
A computer based decision support system and risk chart did not reduce cardiovascular risk or blood pressure in hypertension	Prognosis
Long-acting nifedipine was as effective as hydrochlorothiazide plus amiloride for reducing mortality and morbidity in hypertension	Is chaemic stroke subtypes had different short term and long term functional outcomes, mortality, and recurrence rates $\dots 26$
Diltiazem was as effective as diuretics or β blockers, or both, at preventing cardiovascular mortality and morbidity	Aetiology Review: evidence does not support an association between allergen levels and risk for asthma
Clonidine was effective for reducing tamoxifen associated hot flashes in postmenopausal women with breast cancer 18 Review: palliative chemotherapy reduces death and progression at 12 months in advanced or metastatic colorectal cancer	Clinical prediction guide A simple clinical tool that included age, weight, and oestrogen use helped to select women for bone densitometry
Review: smooth muscle relaxants treat abdominal pain and loperamide reduces diarrhoea in irritable bowel syndrome 20	Glossary Inside back cover