Therapeutics

Formoterol was more effective than terbutaline when taken as needed for moderate to severe asthma

Tattersfield AE, Löfdahl CG, Postma DS, et al. *Comparison of formoterol and terbutaline for as-needed treatment of asthma: a randomised trial*. *Lancet* 2001 Jan 27;357:257–61.

QUESTION: In patients with moderate to severe asthma who use an inhaled corticosteroid but still require as needed medication, is formoterol (a long acting β_2 agonist) more effective than terbutaline (a short acting β_2 agonist) when used as needed?

Design

12 week randomised (allocation concealed*), blinded (patients and investigators initially),* controlled trial.

Setting

35 centres in Greece, the Netherlands, Norway, and Sweden.

Patients

362 patients who were ≥ 18 years of age (mean age 47 y, 57% women); had had asthma for ≥ 6 months; had been treated with a constant dose of an inhaled corticosteroid for ≥ 4 weeks; had an FEV₁ of $\geq 50\%$ of the predicted value, which increased by $\geq 12\%$ after inhalation of 1.5 mg of terbutaline; and used their relief inhaler about 3–8 times per day on ≥ 7 days of the 2 week run in period. Exclusion criteria were need for ≥ 12 inhalations of rescue medication during the run in period or serum potassium level outside of reference range. Follow up was 85%.

Intervention

Patients were allocated to inhaled formoterol, 4.5 μ g (metered dose 6 μ g) (n = 182), or inhaled terbutaline, 0.5 mg (n = 180), for 12 weeks. Patients were told to take the medication only when needed.

Main outcome measures

Time to first severe exacerbation. Secondary outcomes were morning and evening peak flow rate, FEV₁, symptoms, number of inhalations of relief medication, and safety.

Main results

Analysis was by intention to treat. Fewer patients in the formoterol group than in the terbutaline group had ≥ 1 exacerbation (p = 0.02)† (table). The time to first exacerbation was longer in the formoterol group than in the terbutaline group (p = 0.013). Morning and evening peak expiratory flow rates increased in the formoterol group and decreased in the terbutaline group (mean difference 11 l/min, 95% CI 3 to 20 l/min for morning; 8 l/min, CI 0 to 15 l/min for evening). The reduction in number of inhalations of relief medication was higher in the formoterol group than in the terbutaline group (mean difference 0.76 inhalations/d, CI 0.33 to 1.18). Prebronchodilator FEV₁ was increased in the formoterol group relative to the terbutaline group (mean ratio 105%, CI 101% to 108%). Both treatments were well tolerated. Groups did not differ for change in symptom scores.

Conclusion

In patients with moderate to severe asthma, formoterol was more effective than terbutaline when taken as needed.

*See glossary.

†p Value calculated from data in article.

Source of funding:
AstraZeneca R&D.

For correspondence:
Professor A E
Tattersfield, Division of
Respiratory Medicine,
City Hospital,
Nottingham NG5 1PB,
UK. Fax +44 (0)115
840 4771.

Formoterol v terbutaline for moderate to severe asthma†

Outcome at 12 months	Formoterol	Terbutaline	RRR (95% CI)	NNT (CI)
≥ 1 exacerbation	14%	24%	40% (7.6 to 62)	11 (6 to 66)

†Abbreviations defined in glossary; RRR, NNT, and CI calculated from data in article.

COMMENTARY

Formoterol is a long acting, inhaled, β_2 adrenergic receptor agonist with interesting pharmacological properties. Despite having a duration of bronchodilating activity of > 12 hours in patients with asthma, its onset of action is similar to the shorter acting inhaled β_2 adrenergic receptor agonists, such as terbutaline or salbutamol. In addition, the duration of the systemic pharmacological activity of formoterol (resulting in potential side effects) is similar to the shorter acting inhaled β_2 agonists. This characteristic allows formoterol to be used for "as needed" treatment of symptoms.

The study by Tattersfield *et al* compared formoterol and terbutaline used "as needed" in a well designed study in adult patients who had moderately severe and uncontrolled asthma. The main outcome variable, time to first severe asthma exacerbation, is an important outcome in asthma but is not often used in clinical trials.

This study showed that fewer inhalations of "as needed" formoterol were needed and that, somewhat surprisingly, the time to the first severe exacerbation was longer in the formoterol group. This effect on exacerbations has been previously described in a similar patient population, when formoterol, taken regularly twice daily, was added to low or moderate doses of the inhaled corticosteroid budesonide.¹ The current study suggests that the ability of formoterol to reduce the risk for a severe asthma exacerbation is so robust that it can be shown even when the drug is used less frequently. These results are important to clinicians treating asthma because severe asthma exacerbations are the most dangerous events that can occur in patients with asthma, as well as being the most demanding and expensive for the healthcare system. This benefit of formoterol was not accompanied by an increase in β_2 agonist related side effects. The study has convincingly shown that in addition to the already well accepted benefits of the regular use of inhaled formoterol, "as needed" use provides more clinical benefit to people with asthma than does use of the shorter acting terbutaline.

Paul M O'Byrne, MB
McMaster University and St. Joseph's Hospital
Hamilton, Ontario, Canada

1 Pauwels RA, Lofdahl CG, Postma DS, *et al*. Effect of inhaled formoterol and budesonide on exacerbations of asthma. Formoterol and corticosteroids establishing therapy (FACET) international study group. *N Engl J Med* 1997;337:1405–11.