


# Evidence-Based Medicine



**Editors:** Brian Haynes & Paul Glasziou

**Contents**

|  |     |
|--|-----|
| Purpose and procedure  | 130 |
| Additional articles abstracted in <i>ACP Journal Club</i>  | 132 |
| EBM notebook: Creating knowledge management skills in primary care residents: a description of a new pathway to evidence-based practice in the community | 133 |
| Resource corner  | 135 |

**Therapeutics**

|  |     |
|--|-----|
| Review: the levonorgestrel releasing intrauterine system prevents more pregnancies than intrauterine devices (IUDs) $\leq 250$ mm <sup>2</sup> but not IUDs $>250$ mm <sup>2</sup> | 137 |
| Antibiotics did not prevent adverse outcomes after preterm, prelabour rupture of fetal membranes   | 138 |
| Antibiotics did not prevent adverse outcomes in women in preterm labour with intact fetal membranes  | 139 |
| Postexposure prophylaxis with inhaled zanamivir was efficacious for household contacts of people with influenza  | 140 |
| Oseltamivir reduced new cases of influenza in household contacts of people with clinical or confirmed influenza  | 141 |
| Direct observation of tuberculosis treatment did not promote higher cure rates than self administered treatment  | 142 |
| Review: mefloquine prevents malaria but has adverse effects that limit its acceptability in adults not immune to malarial infection  | 143 |
| Incidence of long term backache was similar between women receiving epidural analgesia during labour and those who did not   | 144 |
| Radiography led to improved patient satisfaction but increased short term pain in patients with low back pain  | 145 |
| Review: angiotensin converting enzyme inhibitors are beneficial in type 1 diabetes mellitus and microalbuminuria regardless of baseline risk factor status                         | 146 |

|   |     |
|---|-----|
| Review: oral anticoagulants plus antiplatelet agents reduce thromboembolism and all cause mortality for heart valve prostheses          | 147 |
| A peer led asthma education programme in adolescents was more effective than no programme for improving quality of life                 | 148 |
| Review: some but not all topical antimicrobial agents improve the rate of healing of chronic wounds                                     | 149 |
| Review: warfarin prevents stroke in non-rheumatic atrial fibrillation but has a higher risk for haemorrhage than other agents           | 150 |
| Reduced sodium intake lowered blood pressure and need for antihypertensive medication   | 151 |
| Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for head lice infestation in children | 152 |
| Review: smooth muscle relaxants improve symptoms and reduce pain in irritable bowel syndrome  | 153 |
| A depression management programme reduced depression in frequent users of health care but did not reduce healthcare visits              | 154 |
| A depression management programme increased depression free days and costs in depressed frequent users of general health care           | 155 |
| Patient education to encourage graded exercise improved physical functioning in chronic fatigue syndrome                                | 156 |
| Cognitive behaviour therapy reduced fatigue severity and functional impairment in chronic fatigue syndrome                              | 157 |

**Diagnosis**

|   |     |
|---|-----|
| A normal alveolar dead space fraction plus D-dimer assay ruled out suspected pulmonary embolism in the emergency department | 158 |
| The Mini Cog had high sensitivity and specificity for diagnosing dementia in community dwelling older adults                | 159 |

**Differential diagnosis**

|  |     |
|--|-----|
| A 2 step procedure determined diagnosis in patients with unexplained weight loss | 160 |
|--|-----|