

Evidence-Based Medicine

Contents

Purpose and Procedure	162	Antihypertensive drugs had either a neutral or beneficial effect on lipids	180
Additional Articles Abstracted in <i>ACP Journal Club</i>	163	Service dogs improved the lives of persons with ambulatory disabilities	181
EBM NOTEBOOK		Endarterectomy was not effective for moderate symptomatic carotid stenosis	182
Down with odds ratios!	164	Interferon β -1a slowed progression of disability in multiple sclerosis	183
Research intelligence for policy and practice: the role of the National Health Service Centre for Reviews and Dissemination	167	n of 1 trials led to less theophylline use with no adverse effects	184
Corrections	168	DIAGNOSIS	
THERAPEUTICS		Meta-analysis: Duplex ultrasonography is accurate for most peripheral arterial occlusive disease in legs	185
Meta-analysis: Several strategies prevent falls and subsequent injury in older persons	169	PROGNOSIS	
Elective small-incision cholecystectomy compared with laparoscopy had a shorter operation time and similar recovery	170	Tardive dyskinesia was associated with poor treatment response and higher antipsychotic drug dose after the first episode of schizophrenia	186
Open surgery led to fewer recurrent biliary symptoms than did endoscopic sphincterotomy in high-risk elderly patients	171	ETIOLOGY	
Prophylaxis with ampicillin before surgery was more effective than penicillin after surgery in rural Africa	172	Meta-analysis: Environmental tobacco smoke increases childhood morbidity and mortality	187
Meta-analysis: Calcium supplementation reduces blood pressure and preeclampsia during pregnancy	173	Review: Modest alcohol consumption lowers risk for coronary heart disease irrespective of the type of drink	188
Meta-analysis: Amniotomy does not reduce the rate of caesarean section	174	Inverse association between alcohol intake and risk for ischemic heart disease depended on the LDL cholesterol level	189
Meta-analysis: Prophylactic indomethacin reduces patent ductus arteriosus and intraventricular haemorrhage in premature infants	175	QUALITY IMPROVEMENT	
Pavement stencils marking safe areas promoted safe behaviour at school bus stops	176	Obstetrician-led shared care was not better than care from general practitioners and community midwives in low-risk pregnant women	190
Active compression-decompression CPR for cardiac arrest did not improve survival or neurologic outcomes	177	A reduced schedule of antenatal visits resulted in similar clinical outcomes, poorer psychosocial outcomes, and less satisfaction	191
Meta-analysis: Psychosocial interventions augment cardiac rehabilitation programs	178	ECONOMICS	
Lovastatin was more effective than stepped care in reducing cholesterol	179	The nicotine patch was a cost-effective adjunct to smoking cessation	192
		Glossary	Inside back cover