

Contents

Purpose and procedure	65	Raloxifene reduced vertebral fractures and breast cancer regardless of prior hormone therapy use in women	84
EBM notebook		The cognitive behavioural analysis system of psychotherapy prevented recurrence in chronic major depression	85
Jottings . . .	66	Foot temperature monitoring at home reduced foot complications in high risk patients with diabetes	86
Does clinical experience make up for failure to keep up to date?	66	The addition of peak expiratory flow monitoring to symptom-based self management did not enhance outcome in children with asthma	87
Practice corner: will it happen again doctor? Prognosis after a first seizure	68	Review: bupropion and nortriptyline each increase smoking cessation rates	88
Weighted event rates	70		
Letter: From research evidence to context: the challenge of individualising care	71	Diagnosis	
Evidently . . .	72	Review: a low clinical probability plus a normal D-dimer test result excludes a diagnosis of deep venous thrombosis	89
Therapeutics		Aetiology	
Review: statins reduce stroke but not stroke mortality	73	A haemoglobin concentration of 140 g/l was associated with a lower mortality risk than concentrations ≤ 120 g/l in older women with disabilities	90
Review: atenolol may be ineffective for reducing cardiovascular morbidity or all cause mortality in hypertension	74	Quality improvement	
Diuretic based therapy reduced CV mortality in older patients with isolated systolic hypertension and diabetes	75	A series of evidence-based drug therapy letters improved prescribing behaviour	91
Review: angiotension receptor blockers do not differ from ACE inhibitors in chronic heart failure or acute MI	76	Clinical prediction guide	
Review: ACE inhibitors, but not angiotensin II receptor antagonists, reduce all cause mortality in diabetic nephropathy	77	Self measured home blood pressure was better than "clinic" blood pressure for predicting stroke in a Japanese population	92
Trandolapril did not reduce cardiovascular death or other events in stable coronary artery disease	78	Resource review	93
Picotamide reduced all cause mortality more than aspirin in type 2 diabetes mellitus and peripheral arterial disease	79	Additional articles abstracted in ACP Journal Club	94
Review: low molecular weight heparin reduces recurrent venous thromboembolism better than unfractionated heparin	80	Other articles noted	95
Topical diclofenac improved pain and physical function with no systemic side effects in primary osteoarthritis of the knee	81	Glossary	Inside back cover
Cognitive behaviour therapy reduced relapses in recurrent major depressive disorder	82		
Review: intravenous metoclopramide is better than placebo for reducing pain in acute migraine in the emergency department	83		