

# Evidence-Based Medicine

## for Primary Care and Internal Medicine



[www.evidence-basedmedicine.com](http://www.evidence-basedmedicine.com)

**Editors:** Brian Haynes & Paul Glasziou

### Contents

<b>Purpose and procedure</b>	<b>129</b>	<b>Diagnosis</b>	
<b>EBM notebook</b>		BNP and amino terminal proBNP assays did not differ for detecting congestive heart failure in the emergency department	<b>148</b>
Jottings...	<b>130</b>	A decrease of $\geq 2$ points on the Mini-Mental State Examination was the best determinant of delirium in elderly patients admitted to hospital	<b>149</b>
Letter: Meta-analysis can be statistically misleading	<b>130</b>	<b>Prognosis</b>	
In response: ... but they also present an opportunity to learn more.	<b>130</b>	Review: B type natriuretic peptide consistently predicts death and cardiovascular events in heart failure	<b>150</b>
Clinical skills textbooks fail evidence-based examination	<b>131</b>	Gleason score predicted mortality rate to 20 years for untreated early prostate cancer	<b>151</b>
Statistical approaches to uncertainty: p values and confidence intervals unpacked	<b>133</b>	Overall mortality was 2.2% in children after a first unprovoked afebrile seizure	<b>152</b>
Evidently...	<b>135</b>	<b>Aetiology</b>	
<b>Therapeutics</b>		Subclinical hypothyroidism increased the risk of placental abruption and poor neonatal outcomes	<b>153</b>
Warfarin was not more effective than aspirin and increased adverse events in symptomatic intracranial arterial stenosis	<b>136</b>	<b>Quality improvement</b>	
Low dose aspirin lowered stroke risk but not risks of MI or cardiovascular deaths in women	<b>137</b>	Pharmacist led, primary care based disease management reduced risk factors and improved glycaemic control in diabetes	<b>154</b>
Intensified blood pressure (BP) control was not better than conventional BP control in non-diabetic chronic renal disease	<b>138</b>	<b>Clinical prediction guide</b>	
ED use of magnesium sulphate improved rate control in atrial fibrillation with rapid ventricular response	<b>139</b>	A clinical prediction model predicted absence of significant fibrosis in chronic hepatitis B	<b>155</b>
Review: several pharmacological therapies promote modest weight loss	<b>140</b>	<b>Additional articles abstracted in ACP Journal Club</b>	<b>156</b>
Review: sparse high quality evidence supports surgery for obesity	<b>141</b>	<b>Other articles noted</b>	<b>157</b>
Screening for abdominal aortic aneurysm (AAA) reduced AAA mortality in Danish men 64–73 years of age	<b>142</b>	<b>Resource reviews</b>	<b>159</b>
A smoking cessation programme plus inhaled ipratropium improved survival in asymptomatic airway obstruction	<b>143</b>	<b>Glossary</b>	Inside back cover
Clinician advice, an interactive computer program, and motivational counselling increased smoking cessation in teens	<b>144</b>		
Cognitive therapy reduced relapses and number of days in bipolar episodes in DSM-IV bipolar I disorder	<b>145</b>		
Review: bright light therapy and dawn simulation reduce symptom severity in seasonal affective disorder	<b>146</b>		
Buccal oxycodone reduced pain more than placebo in children with acute undifferentiated abdominal pain	<b>147</b>		