

# Evidence-Based Medicine

## for Primary Care and Internal Medicine



[www.evidence-basedmedicine.com](http://www.evidence-basedmedicine.com)

**Editors:** Brian Haynes & Paul Glasziou

### Contents

|  |           |   |                   |
|--|-----------|---|-------------------|
| <b>Purpose and procedure</b>   | <b>1</b>  | Nurse-led cognitive behaviour therapy improved insight and reduced negative symptoms and readmissions in schizophrenia                | <b>21</b>         |
| <b>EBM notebook</b>  |           | Periodic telephone counselling for polypharmacy improved compliance and reduced mortality   | <b>22</b>         |
| Jottings...  | <b>2</b>  | Review: opiate administration may alter physical examination findings but does not increase management errors in acute abdominal pain | <b>23</b>         |
| Misunderstandings, misperceptions, and mistakes  | <b>2</b>  |   |                   |
| The evolving science of translating research evidence into clinical practice   | <b>4</b>  |   |                   |
| <b>Evidently...</b>  | <b>8</b>  | <b>Diagnosis</b>  |                   |
| <b>Therapeutics</b>  |           | Review: tissue transglutaminase test is almost as accurate as endomysial antibody test for diagnosis of coeliac disease               | <b>24</b>         |
| Budesonide-formoterol for maintenance and as needed reliever treatment reduced asthma exacerbations                                  | <b>9</b>  | Review: history and physical examination can accurately identify migraine and the need for neuroimaging in patients with headache     | <b>25</b>         |
| Review: long acting $\beta$ agonists increase severe asthma exacerbations and asthma related deaths in children and adults           | <b>10</b> | <b>Aetiology</b>  |                   |
| Review: omalizumab reduces exacerbation and steroid use in chronic asthma  | <b>11</b> | Fetal exposure to ACE inhibitors increased risk of major congenital malformations   | <b>26</b>         |
| Review: long acting $\beta_2$ adrenoceptor agonists are effective in poorly reversible chronic obstructive pulmonary disease         | <b>12</b> | <b>Quality improvement</b>  |                   |
| Review: anticholinergics but not $\beta_2$ agonists reduce exacerbations requiring hospital admission and respiratory deaths in COPD | <b>13</b> | Review: 3 of 4 RCTs on the treatment of adolescent depression in primary care have positive results                                   | <b>27</b>         |
| Physiotherapy plus a craniocervical training programme was better than physiotherapy alone in tension type headache                  | <b>14</b> | <b>Economics</b>  |                   |
| Review: medical therapy with calcium channel blockers or $\alpha$ blockers helps patients to pass urinary stones                     | <b>15</b> | B type natriuretic peptide testing was more cost effective than conventional diagnosis in patients with acute dyspnoea                | <b>28</b>         |
| Review: planned early birth after prelabour rupture of membranes at term has benefits for mother and infant                          | <b>16</b> | <b>Resource review</b>  | <b>29</b>         |
| Diuretics were superior to calcium channel blockers and short term ACE inhibitors for reducing heart failure in hypertension         | <b>17</b> | <b>Additional articles abstracted in ACP Journal Club</b>   | <b>30</b>         |
| Daily oral vitamin B-12 with or without folic acid for 24 weeks did not improve cognitive function in elderly people                 | <b>18</b> | <b>Other articles noted</b>   | <b>31</b>         |
| Environmental and dietary interventions in the first 5 years of life did not reduce risk of asthma and allergic disease              | <b>19</b> | <b>Glossary</b>   | Inside back cover |
| Naltrexone increased abstinence and reduced heavy drinking during treatment  | <b>20</b> |   |                   |