

Evidence-Based Medicine

for Primary Care and Internal Medicine

Editors: Brian Haynes & Paul Glasziou



www.evidence-basedmedicine.com

Contents

Purpose and procedure	97	Once daily oral controller therapy with low dose theophylline or montelukast was not effective in poorly controlled asthma	115
EBM notebook		Review: drugs for mild to moderate hypertension in pregnancy reduce the risk of severe hypertension but not pre-eclampsia	116
Jottings...	98	Reducing infant exposure to food and dust mite allergens reduced the incidence of asthma and allergy at age 8 years	117
The 6 steps of evidence-based medicine	98	Review: periodic health examination increases delivery of some clinical preventive services and reduces patient worry	118
6Ts teaching tips for evidence-based practitioners	100	Stress echocardiography safely classified more patients as low risk of serious CAD than exercise electrocardiography	119
Adding cost to number needed to treat	101		
Letter	102		
Evidently...	103		
Therapeutics		Diagnosis	
Circumcision reduced the risk of contracting HIV infection in young sexually active Kenyan men	104	Review: ultra-short screening tests are not highly accurate for detecting depression in primary care	120
Review: flight times >8 hours and presence of risk factors for VTE increases travel related VTE	105	Review: heterogeneous studies show that nucleic acid amplification tests accurately rule in but do not rule out TB	121
Intra-articular corticosteroid injections with the reciprocating procedure device reduced procedural pain and duration more than the conventional syringe	106	Clinical prediction guide	
Percutaneous coronary intervention plus optimal medical therapy was not more effective than medical therapy alone in stable CAD	107	The 4 item Fracture and Mortality Index predicted hip fracture and all cause mortality in elderly women	122
Review: lifestyle or pharmacological interventions prevent or delay type 2 diabetes in impaired glucose tolerance	108	Prognosis	
Review: collaborative care was effective for depression in primary care in the short and longer term	109	Absolute risk of subsequent fracture was similar in women and men	123
A hospital-based violence prevention intervention reduced hospital recidivism for violent injury and arrests for violent crimes	110	Resource review	124
Booklet-based education in vestibular rehabilitation or symptom control improved subjective health in Ménière disease	111	Additional articles abstracted in ACP Journal Club	126
Review: available evidence does not support the use of β blockers as first line treatment for hypertension	112	Other articles noted	127
An intensive smoking cessation intervention reduced hospital admissions and mortality in high risk smokers with CVD	113	Glossary	Inside back cover
Salmeterol and fluticasone propionate did not reduce mortality in chronic obstructive pulmonary disease	114		