

Evidence-Based Medicine

for Primary Care and Internal Medicine

Editors: Brian Haynes & Paul Glasziou



www.evidence-basedmedicine.com

Contents

Purpose and procedure	129		
EBM notebook			
Jottings . . .	130		
Putting evidence into context: some advice for guideline writers	130		
Practice corner: Chair's rounds	133		
Evidently	135		
Therapeutics			
Long term use of eicosapentaenoic acid reduced major coronary events in hypercholesterolaemia	136		
Hormone therapy for younger women may not increase CHD risk during 5–7 years follow-up, but stroke risk was increased independent of age	137		
The Atkins diet led to more weight loss than the Zone diet in overweight and obese premenopausal women at 12 months	138		
A care management intervention improved depression after stroke	139		
A self-management approach to patient education for type 2 diabetes was more effective than a didactic approach	140		
Carvedilol was more effective than metoprolol for preventing cardiovascular events in heart failure	141		
An escalating higher energy regimen was better than a fixed lower energy regimen for defibrillation in out of hospital cardiac arrest	142		
6 or 3 months of anticoagulant therapy did not differ for treatment failure in patients with DVT, PE, or both	143		
Ibuprofen was more effective than codeine or acetaminophen for musculoskeletal pain in children	144		
A once yearly IV infusion of zoledronic acid prevented fractures in postmenopausal women with osteoporosis	145		
10 day sequential therapy was more effective than 10 day triple drug therapy for eradicating <i>Helicobacter pylori</i> infection	146		
Review: partner notification interventions can reduce sexually transmitted infections	147		
		Ambulatory titration of continuous positive airway pressure was as effective as polysomnography for obstructive sleep apnoea	148
		Diagnosis	
		The 2-item Generalized Anxiety Disorder scale had high sensitivity and specificity for detecting GAD in primary care	149
		A short version of Mini-Mental State Examination (MMSE) was as accurate as the original MMSE for predicting dementia	150
		Review: clinical examination is often as accurate as magnetic resonance imaging for diagnosing meniscus tears	151
		Review: glycated haemoglobin A _{1c} and fasting plasma glucose screening tests have similar sensitivities and specificities for early detection of type 2 diabetes	152
		Economics	
		Routine rotavirus vaccinations of infants cost US\$138 per case averted and US\$197 190 per life year saved from a societal perspective	153
		Clinical prediction guide	
		A clinical prediction guide predicted progression to rheumatoid arthritis in undifferentiated arthritis	154
		Pre-endoscopic serological test with duodenal biopsy in high risk patients had high sensitivity and low specificity for coeliac disease	155
		Resource review	156
		Additional articles abstracted in ACP Journal Club	158
		Other articles noted	159
		Glossary	Inside back cover