

Evidence-Based Medicine

for Primary Care and Internal Medicine

Editors: Brian Haynes & Paul Glasziou



www.evidence-basedmedicine.com

Contents

Purpose and procedure	161	Review: brief interventions including information sessions, motivational interviews, and cognitive behavioural therapy reduce excessive alcohol consumption in primary care	179
EBM notebook		An intensive lifestyle intervention reduced weight and cardiovascular disease risk factors in obese participants with type 2 diabetes at 1 year in an interim analysis	180
Jottings . . .	162	Irrigation of simple lacerations with tap water or sterile saline in the emergency department did not differ for wound infections	181
Life after the EBM workshop	163		
Smart health choice essentials	164		
Letter	166		
Evidently	167		
Therapeutics		Diagnosis	
Review: vaccines against human papillomavirus prevent cervical cancer precursors in young women for at least 3 years	168	Review: some screening tests for dementia in older people are accurate and practical for use in primary care	182
Review: rosiglitazone increases risk of MI but does not differ from other drugs for CV death in type 2 diabetes	169	Review: anti-cyclic citrullinated peptide antibody is a more specific test for rheumatoid arthritis than rheumatoid factor	183
Rosiglitazone increased heart failure but did not differ from metformin plus sulphonylurea for other CV outcomes at interim analysis	170	Aetiology	
Review: regular brisk walking improves cardiovascular risk factors in healthy sedentary adults	171	Review: increased waist circumference or waist-to-hip ratio is associated with increased risk of cardiovascular events	184
Warfarin reduced major stroke more than aspirin in elderly patients with atrial fibrillation in primary care	172	Quality improvement	
Symptomatic use of beclomethasone-albuterol and regular use of beclomethasone did not differ for control of mild asthma	173	Reminders in echocardiography reports increased use of β blockers in reduced left ventricular ejection fraction	185
Induction of labour post-term or expectant management with serial fetal monitoring had similar neonatal outcomes	174	Clinical prediction guide	
Prednisolone plus paracetamol (acetaminophen) was as effective as indomethacin plus paracetamol but had fewer adverse effects in acute gout-like arthritis	175	Review: evidence from single studies shows that a few fall risk assessment tools can predict falls in elderly people	186
Intranasal fentanyl and intravenous morphine did not differ for pain relief in children with closed long-bone fractures	176	The ankle brachial index independently predicted all-cause mortality in high-risk patients with peripheral arterial disease	187
Review: on-demand maintenance therapy with proton pump inhibitors is as effective as continuous therapy for non-erosive GORD	177	Resource review	188, 189
Cleaning of the perineal/genital area before urine collection from toilet-trained children prevented sample contamination	178	Additional articles abstracted in ACP Journal Club	190
		Other articles noted	191
		Glossary	Inside back cover