


# Evidence-Based Medicine



## Contents

April 2008 Vol 13 No 2

### Purpose and procedure

33

### EBM notebook

- 34 Jottings ...
- 34 Epley and the slow boat from research to practice
- 36 Better value—the 21st century priority

### Evidently

39

### Therapeutics

- 40  Aerobic plus resistance training was more effective than either alone for reducing HbA<sub>1c</sub> levels in type 2 diabetes
- 41 Review: diet interventions, with or without exercise, promote weight loss more than advice alone
- 42 Review: delayed or immediate prescriptions of antibiotics have similar clinical outcomes in respiratory infections
- 43 Both low-dose and micro-dose 17 $\beta$ -oestradiol reduced hot flushes
- 44 Early treatment with prednisolone, but not acyclovir, was effective in Bell's palsy
- 45 Systematic or opportunistic screening was more effective than usual care for detecting new cases of atrial fibrillation
- 46 A probiotic drink prevented diarrhoea and *Clostridium difficile* infection in older patients taking antibiotics
- 47 Review: vitamin D supplementation decreases all-cause mortality in adults and older people
- 48 Daily vitamin C prevented development of complex regional pain syndrome in adults with wrist fractures
- 49 Perindopril plus indapamide reduced death and major macrovascular and microvascular events in type 2 diabetes
- 50 Review:  $\beta$ -blockers for hypertension increase risk of new onset diabetes

- 51 Treatment review by case conferences led to more medication changes than written feedback in older people on polypharmacy
- 52 Combined HRT increased CV events more than placebo but did not differ from oestrogen HRT after menopause
- 53 Review: calcium supplementation, with or without vitamin D, prevents osteoporotic fractures in older people
- 54 A 16-week tai chi programme prevented falls in healthy older adults
- 55 Pharmacogenetic-guided and standard dosing did not differ for out-of-range INRs in patients initiating warfarin therapy
- 56 Review: long-acting inhaled therapies and pulmonary rehabilitation are effective in stable COPD
- 57 Review: corticosteroids plus long-acting  $\beta$ -agonists reduce exacerbations more than long-acting  $\beta$ -agonists alone in COPD

### Diagnosis

- 58 Accuracy of ECG interpretation in primary care was limited for detecting atrial fibrillation

### Quality improvement

- 59 A composite management intervention plus financial incentives reduced mortality from malaria in children <5 years of age

### Resource reviews

60

### Additional articles abstracted in ACP Journal Club

62

### Other articles noted

63

### Glossary

Inside back cover

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.



[www.publicationethics.org.uk](http://www.publicationethics.org.uk)  
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics