


# Evidence-Based Medicine



## Contents

June 2008 Vol 13 No 3

### Purpose and procedure

65

### EBM notebook

- 66 Practice corner: managing trichotillomania—no need to pull your hair out
- 67 Interpreting meta-analysis in systematic reviews


### Letter

- 69 Proactive monitoring for newly emerging evidence: The lost step of EBP?

### Evidently

70

### Therapeutics

- 71 Testosterone supplementation did not prevent cognitive decline or increase bone mineral density in older men
- 72  Review: use of pedometers increases physical activity in adults
- 73 Review: combination therapy with renin angiotensin inhibitors reduces proteinuria more than single drugs alone in renal disease
- 74 Review: corticosteroids plus LABAs reduce exacerbations and mortality more than steroids alone in COPD
- 75 Adjunctive diclofenac and spinal manipulation did not speed recovery of acute low back pain
- 76 Review: individual DMARDs have similar efficacy for RA, but combination therapy improves response
- 77 Adjunctive prandial or biphasic insulin for type 2 diabetes improved glycaemic control but increased hypoglycaemia and weight gain
- 78 A patient-adjusted insulin algorithm was non-inferior to standard of care for glycaemic control in type 2 diabetes
- 79 Review: long-acting insulin analogues do not improve glycaemic control but reduce nocturnal hypoglycaemia
- 80 5 days of nitrofurantoin was equivalent to 3 days of trimethoprim-sulfamethoxazole for women with non-complicated cystitis

- 81 Review: multiple doses of antenatal corticosteroids prevent preterm neonatal respiratory disease more than a single dose
- 82 Review: misoprostol and intramuscular prostaglandins do not prevent postpartum haemorrhage more than injectable uterotonics
- 83 Review: calcium supplementation during pregnancy reduces the risk of pre-eclampsia

### Diagnosis

- 84 Random urine protein:creatinine ratio was an accurate method for diagnosing proteinuria in pregnant women with hypertension
- 85 Review: non-psychiatric physicians have low accuracy for recognising depression in their patients

### Prognosis

- 86 Review: risk of stroke is high at 2, 30, and 90 days after transient ischaemic attack

### Aetiology

- 87 Review: lower total cholesterol is associated with reduced risk of death from ischaemic heart disease but not stroke
- 88 Review: silent brain infarcts are common in the elderly general population
- 89 Glomerular filtration rate  $< 45$  ml/min/1.73 m<sup>2</sup> with microalbuminuria was associated with increased CV death
- 90 BMI and cardiorespiratory fitness predicted mortality in older adults

### Resource reviews


92

### Other articles noted

95

### Glossary

Inside back cover

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 [www.publicationethics.org.uk](http://www.publicationethics.org.uk)  
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics