

Evidence-Based Medicine



Contents

October 2008 Vol 13 No 5

Purpose and procedure

129

EBM notebook

130 Jottings ...

130 Evidence-based medicine targets the individual patient, part 2: guides and tools for individual decision-making


132 Spectrum bias: why generalists and specialists do not connect

133 Why A-level evidence does not make it to clinicians' A-list: the case of thromboprophylaxis in medical patients

Evidently

135

Therapeutics

136  Antihypertensive therapy with indapamide and perindopril reduced mortality in patients ≥ 80 years

137 A list of 10 tips improved weight control in overweight and obese adults

138 Review: special diets, including egg and cow's milk exclusion, a few-foods diet, and an elemental diet, are not effective in unselected cases of atopic eczema

139 Review: little high-quality evidence is available regarding treatments for localised prostate cancer

140 Review: limited evidence suggests that statins are safe in patients with liver disease

141 Buflomedil reduced cardiovascular events in peripheral arterial disease

142 Risk of cancer was still increased over 2 years after stopping hormone therapy

143 Imiquimod 5% cream reduced lesion size in vulvar intraepithelial neoplasia

144 Laparoscopic appendicectomy reduced chronic right lower-quadrant abdominal pain more than sham intervention

145 Oral prednisolone was not inferior to intravenous prednisolone for treatment failure in chronic obstructive pulmonary disease exacerbation

146 Metformin plus a lifestyle intervention was more effective than either alone for antipsychotic-induced weight gain

147 Telmisartan and ramipril were equivalent, but their combination increased adverse events in vascular disease or diabetes

148 Review: some evidence supports pharmacological and non-pharmacological treatments of fibromyalgia syndrome

149 Review: erythropoiesis-stimulating agents increase mortality and venous thromboembolism in cancer-associated anaemia

150 Metformin (alone or with insulin) was as effective as insulin for preventing perinatal complications in gestational diabetes

Aetiology

151 Non-diabetic maternal hyperglycaemia was associated with adverse pregnancy outcomes

Diagnosis

152 Review: simple questions and clinical tests are moderately useful for diagnosing urinary incontinence

Prognosis

153 HF patients with preserved or reduced ejection fractions did not differ for survival or CV mortality at 5 years

154 Risk of fatal pulmonary embolism was 0.49 per 100 person-years after discontinuing anticoagulant therapy for venous thromboembolism

155 Melanoma of the scalp and neck had greater risk of melanoma-specific mortality than melanoma of the extremities

Resource reviews

156

Other articles noted

159

Glossary

Inside back cover



This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk