


# Evidence-Based Medicine



## Contents

December 2008 Vol 13 No 6

### Purpose and procedure

161

### EBM notebook

- 162 Jottings ...
- 162 GRADE: assessing the quality of evidence for diagnostic recommendations
- 164 Applying evidence: what's the next action?
- 166 A simple paper-based patient decision aid

### Evidently

167

### Therapeutics

- 168 Intensive glucose control did not prevent important complications in type 2 diabetes
- 169 Intensive glucose control increased mortality and did not prevent cardiovascular events in type 2 diabetes
- 170 Glycaemic control was not affected by self-monitoring of blood glucose in type 2 diabetes
- 171 Self-monitoring of blood glucose was not cost-effective in non-insulin-treated type 2 diabetes
- 172 Review: little evidence exists on the benefits and harms of screening for and treating gestational diabetes
- 173 Lifestyle interventions reduced the long-term risk of diabetes in adults with impaired glucose tolerance
- 174 Review: BP-lowering drugs prevent cardiovascular events equally well in younger and older patients
- 175 Metoprolol prevented myocardial infarction but increased risk of stroke and death after non-cardiac surgery
- 176 Low-carbohydrate and Mediterranean diets led to greater weight loss than a low-fat diet in moderately obese adults
- 177 Review: antioxidant supplements do not reduce all-cause mortality in primary or secondary prevention

- 178 Review: antiviral prophylaxis in pregnant women reduces recurrence of genital herpes at delivery
- 179 Review: infection rates do not differ for wounds cleansed with water or saline
- 180 Placebo acupuncture improved symptoms and quality of life in irritable bowel syndrome
- 181 An intervention promoting exclusive and prolonged breast feeding improved verbal intelligence scores in children at 6.5 years
- 182 Early, abrupt weaning did not reduce risk of HIV infection or mortality in breast-fed infants of mothers with HIV
- 183 High-concentration capsaicin reduced pain in patients with HIV-associated polyneuropathy .
- 184 Methylnaltrexone reduced opioid-induced constipation in patients with terminal illness
- 185 Early surgery was better than conservative care for short-term disability and pain in sciatica

### Aetiology

- 186 Higher values of fasting plasma glucose within the normal range were associated with increased risk of type 2 diabetes

### Prognosis

- 187 Review: many adults still have pain and subjective instability at 1 year after acute lateral ankle sprain

### Resource reviews

188

### Additional articles abstracted in ACP Journal Club

190

### Other articles noted

191

### Glossary

Inside back cover

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org.uk](http://www.publicationethics.org.uk)