

Evidence-Based Medicine



Contents

Editorial

- 165** Multimorbidity: a challenge for evidence-based medicine



Therapeutics

- 167** Long-acting β -agonist step-up therapy is more likely to provide best response, compared to inhaled corticosteroid or leukotriene-receptor antagonist step-up in children with uncontrolled asthma receiving inhaled corticosteroids

- 168** Meta analysis suggests that folic acid supplementation does not reduce risk of stroke, but there may be some benefit when given in combination with vitamins B6 and B12 and in primary prevention

- 170** Low-dose and high-dose colchicine have comparable efficacy in the treatment of acute gout, but high dose carries significantly greater risk of adverse effects

- 171** Moderate quality evidence that compared to advice to rest in bed, advice to remain active provides small improvements in pain and functional status in people with acute low back pain

- 172** Sufficient evidence that 12 g of lactose is tolerated by most adults with lactose malabsorption and intolerance but insufficient evidence on the effectiveness of therapeutic strategies tested so far

- 174** Fibrates reduce the risk of major cardiovascular and coronary events compared with placebo, but do not affect risk of cardiovascular or all-cause mortality

- 175** Administration of annual oral high-dose vitamin D to community-dwelling older women in autumn and winter months increases risk of falls and fractures

- 177** Topical NSAIDs provide effective relief of acute musculoskeletal pain compared to placebo, with no increase in risk of adverse effects

- 178** No current evidence that routine administration of oxygen to people with acute myocardial infarction improves pain or mortality; further conclusive trials are needed

- 179** Intensive dietary advice significantly improves HbA(1c) in patients with type II diabetes who remain hyperglycaemic despite optimised drug treatment

December 2010 Vol 15 No 6

- 181** Calcium supplements increase risk of myocardial infarction

Diagnosis

- 182** Single HPV test not useful for predicting CIN2 or worse or for guiding choice of further investigations for women aged 20-59 presenting to NHS Cervical Screening Programme with borderline abnormalities or mild dyskaryosis

- 183** Specific clinical findings, including coma, neck stiffness and seizures, increase the likelihood of haemorrhagic stroke, but no combination of features is definitively diagnostic

Prognosis

- 185** Intraoperative transfusion in older patients with low preoperative haematocrit levels and substantial blood loss during major non-cardiac surgery is associated with lower 30-day postoperative mortality

- 186** About half of children under age 3 whose parents suspected acute otitis media do not have the diagnosis; restless sleep, ear rubbing, crying, irritability and fever are not predictive

- 187** Statin use associated with increased risk of cataract, myopathy, liver dysfunction and acute renal failure with varying numbers needed to harm

Aetiology

- 189** High blood pressure while taking antithrombotic medication is associated with an increased risk of developing intracranial haemorrhage

- 190** Use of oestrogen only hormone replacement therapy associated with increased risk of asthma onset in postmenopausal women

- 191** Risk of neonatal death is increased by birth outside the normal working week for singleton infants with cephalic presentation born at term

- 193** Meta-analysis of results from case control and cohort studies finds that migraine is associated with approximately twice the risk of ischaemic stroke

Electronic pages

- e1** Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

 equator network

 recycle

When you have finished with this please recycle it