


Evidence-Based Medicine



Contents

Editorial

- 161** The intuitive appeal of case series thinking: a challenge for evidence-based teaching and practice

Methods

- 163** A qualitative approach to Bayes' theorem



Aetiology

- 168** Older women who use bisphosphonate for longer than 5 years may have increased odds of a subtrochanteric or femoral shaft fracture, but absolute risk is low



- 169** People aged above 65 treated for latent tuberculosis are at increased risk of admission for a hepatic event compared with untreated controls

- 170** Exposure to diagnostic radiation and risk of childhood cancer: overstated risks raise unnecessary concern

Diagnosis

- 172** Automated blood pressure readings in primary care demonstrate better correlation to the gold standard of ambulatory monitoring than manual assessment

- 173** Two sputum samples at once for diagnosis of tuberculosis in Africa has equivalent sensitivity and specificity to the standard initial sample followed by a morning sample strategy

- 174** A single Xpert MTB/RIF test of sputum for diagnosis of tuberculosis and multidrug resistance shows high sensitivity and specificity and reduces diagnosis and treatment delays

Prognosis

- 176** Physicians' calling patients on excess weight may provide reality check and increase desire to lose weight in overweight and obese individuals

- 177** Introduction of solids before 4 months is associated with obesity at 3 years among formula-fed infants but not among breast-fed infants

December 2011 Vol 16 No 6

- 178** School-aged children who were exposed to sodium valproate in utero have impaired language scores when compared with a population mean score

Therapeutics

- 180** Oral misoprostol reduces the risk of postpartum haemorrhage in home births assisted by trained traditional birth attendants in Pakistan

- 181** Acute otitis media: antibiotics are moderately effective and mildly increase the risk of adverse effects; prevalence of different causative bacteria changed after introduction of the heptavalent pneumococcal conjugate vaccine

- 182** Continuous support in labour has beneficial effects for mother and baby

- 183** In children and adolescents with mild persistent asthma, daily beclomethasone reduces treatment failure compared with rescue beclomethasone plus albuterol

- 185** Sustained reduction in body mass index and other cardiovascular risk markers a year after a 12-month intensive family-based lifestyle intervention for obese children; but follow-up of participants low

- 186** Intensive glucose lowering (HbA1c target <6.0%) for people with type 2 diabetes increases mortality

- 187** In people with atrial fibrillation unsuitable for warfarin, apixaban reduces the risk of stroke compared with aspirin, with no difference in major bleeding

- 189** Tiotropium mist inhaler for COPD increases risk of mortality compared with placebo

- 190** Episodic physical and sexual activity increase risk for acute cardiac events

Letters

- 192** Conflicts of interest

- 192** Editor's reply to 'conflicts of interest'

Electronic page

- e3** Purpose and procedure



This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator network

recycle

When you have finished with this please recycle it