


Evidence-Based Medicine



Contents

June 2012 Vol 17 No 3

Editorial

- 69** For ethical, economic and scientific reasons, health-relevant degree theses must be made publicly accessible



Original EBM research

- 71** How much doctoral research on clinical topics is published?

Systematic review

- 75** Mechanisms of reducing postoperative pain, nausea and vomiting: a systematic review of current techniques

Therapeutics

- 81** High-quality evidence that spinal manipulative therapy for chronic low back pain has a small, short-term greater effect on pain and functional status compared with other interventions
- 82** Averaging clinic and home measures associated with reductions in within-patient variability and may give a better indication of blood pressure control
- 83** Oxytocin for slow progress in the first stage of spontaneous labour is not associated with different rates of caesarean or assisted delivery compared with no treatment or delayed use in second stage
- 85** In people with newly diagnosed type 2 diabetes an intensive dietary intervention, with or without an activity programme, improves glycaemic control over 12 months compared with usual care
- 86** Early supplemental parenteral nutrition in critically ill adults increased infections, ICU length of stay and cost
- 87** Uterine artery embolisation associated with greater need for reintervention than surgical treatment for symptomatic uterine fibroids; quality of life similar though study underpowered

- 88** Metformin is not significantly different from insulin for preventing fetal macrosomia in women with gestational diabetes
- 90** Inhaled corticosteroids for people with chronic obstructive pulmonary disease are associated with an increased risk of fracture
- 91** In premenopausal women with recurrent UTI, antibiotics were more effective than capsules of cranberry extract for the prevention of UTI, but resistance was higher in the antibiotic group
- 92** Review: excisional, and perhaps ablative treatments for precancerous cervical changes increase the risk for preterm birth
- 93** Tiotropium is superior to salmeterol in reducing frequency of exacerbations: but the effect of adding tiotropium to the combination of inhaled corticosteroid and long-acting $\beta(2)$ -agonist remains unclear
- 95** Antiretroviral combination therapy markedly reduces risk of heterosexual HIV-1 transmission
- 96** Varenicline for 4 weeks prior to target quit date reduces prequit date smoking and increases 12-week abstinence
- 98** Intensive glucose-lowering therapy in people with type 2 diabetes: what do we learn from a new meta-analysis of randomised controlled trials?

Prognosis

- 101** Stopping smoking in the weeks prior to surgery has no effect on the risk of postoperative complications

Electronic page

- e3** Purpose and procedure



This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator network

recycle

When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

