

Evidence-Based Medicine



Contents

August 2012 Vol 17 No 4

Editorial

- 103** The 11th hour – time for *EBM* to return to first principles?



Perspective

- 105** Payer decision-making with limited comparative and cost effectiveness data: the case of new pharmacological treatments for gout

Therapeutics

- 109** Robustness, risk and responsivity in early language acquisition: a randomised trial of a low-intensity language promotion programme for slow to talk toddlers finds no effects on language or behavioural development
- 110** Caudal epidural steroid injections no better than saline epidurals or sham injections for the treatment of chronic lumbar radiculopathy
- 111** Some oral antidiabetic medications may prevent Type 2 diabetes in people at high risk for it, but the evidence is insufficient to determine whether the potential benefits outweigh the possible risks
- 112** Oral ondansetron decreases vomiting, as well as the need for intravenous fluids and hospital admission, in children with acute gastroenteritis
- 114** In irritable bowel syndrome, antispasmodics and antidepressants improve abdominal pain and global assessment and symptom scores, but there is no evidence for the effectiveness of bulking agents
- 115** In TIA or stroke patients with intracranial arterial stenosis, aggressive medical therapy was superior to percutaneous transluminal angioplasty and stenting for 30-day risk of further stroke or death
- 116** Evening dosing of antihypertensive medications results in better blood pressure control and decreases cardiovascular morbidity and mortality in patients with Type 2 diabetes

Prevention

- 118** Population benefits of HPV vaccination for boys: a complex equation

Prognosis

- 120** Greater diabetes-care contact frequency is associated with more rapid achievement of care goals but interpretation of finding requires consideration of patient adherence and motivation

- 121** Preterm birth, a known risk factor for infant and childhood death, is an independent risk factor for mortality in early childhood and young adulthood

- 122** Cardiovascular risks of stopping antiplatelet therapy before non-cardiac surgery for patients with coronary stents

Aetiology

- 124** Early introduction of allergenic food is not associated with increased report of wheeze or eczema
- 125** Breastfeeding by mothers infected with hepatitis B carries no increased risk of transmission to infants who receive proper immunoprophylaxis: a meta-analysis
- 126** Breast feeding is associated with decreased risk of sudden infant death syndrome
- 127** Positive parenting in adolescence is associated with well-(adaptive-) functioning adult offspring
- 128** Observational studies find association between chocolate consumption and reduced risk of cardiovascular disease and diabetes

Harm

- 130** Exposure to trimethoprim–sulfamethoxazole is associated with hospitalisation for hyperkalaemia in older people treated with spironolactone

Diagnosis


- 133** Non-fasting blood testing for lipid screening in children result in statistically significant, but not clinically significant, changes in lipid levels

PostScript

- 134** Correction

About EBM

- e4** Purpose and procedure

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it