

Evidence-Based Medicine



Contents

Perspective

- 161** Opinion editorials: the science and art of combining evidence with opinion



Methods

- 165** Individual patient data meta-analysis of diagnostic studies: opportunities and challenges

Original EBM research

- 170** How to confidently teach EBM on foot: development and evaluation of a web-based e-learning course

Therapeutics

- 173** Montelukast potentially efficacious in children with non-severe obstructive sleep apnoea in the short term
- 174** Topical NSAIDs provide effective pain relief for patients with hand or knee osteoarthritis with similar efficacy, and fewer side effects, than oral NSAIDs
- 175** Statins in patients with CKD prove beneficial in reducing cardiovascular events and mortality but show no benefit in patients on dialysis
- 176** While hormone therapy unlikely harmful in younger postmenopausal women at low CV risk, protective effects cannot be claimed
- 177** Opioid substitution therapy is associated with decreased HIV transmission among people who inject drugs
- 179** Tiotropium bromide triple combination therapy improves lung function and decreases asthma exacerbations
- 180** Unclear whether energy-based vessel sealing in vaginal hysterectomy is superior to conventional clamping and suturing
- 181** Use of cranberry products does not appear to be associated with a significant reduction in incidence of recurrent urinary tract infections
- 182** Network meta-analysis shows that prostaglandin inhibitors and nifedipine are best short-term tocolytics for preterm delivery

Prevention

- 184** Prophylactic antiretroviral HIV therapy prevents infection in heterosexual men and women
- 185** Children gain less weight and accumulate less fat when sugar-free, non-caloric beverages are substituted for sugar-sweetened beverages

Diagnosis

- 187** Pretest probability assessment combined with point-of-care D-dimer testing allows primary care physicians to rule out pulmonary embolism

Prognosis

- 189** Need remains for evidence-based reassurance for absence of bilirubin neurotoxicity with severe hyperbilirubinaemia

October 2013 Volume 18 No 5

- 190** Women whose cervical intraepithelial neoplasia has been treated or resolved remain at continued high risk for cervical cancer

Aetiology

- 192** Proton pump inhibitors do not induce change in bone mineral density in a long-term observational study
- 193** Proton pump inhibitor use may be associated with an increased risk of *Clostridium difficile* infection
- 194** The possible effects of depressive symptoms on risk of preterm birth are clouded by lack of control over confounding factors
- 195** It is still an open question whether inhaled glucocorticoid-induced effects may reduce adult height

Quality improvement

- 197** Ambulatory test result follow-up represents a serious problem and solutions are needed

Economic analysis

- 199** After a negative screening colonoscopy, a microsimulation model shows that currently recommended strategies are equally effective for rescreening

Online articles

- e41** Antibiotics are not beneficial for patients with clinically diagnosed uncomplicated acute rhinosinusitis
- e42** Exenatide may improve maintenance of HbA1c targets, with less hypoglycaemia, but more early adverse effects when compared with low-dose glimepiride
- e43** Vaccination of older adults against herpes zoster is safe and effective
- e44** Screening for type 2 diabetes does not reduce mortality over 10 years
- e45** Currently available smartphone apps for asthma have worrying deficiencies
- e46** Nucleic acid amplification tests of self-taken vulvovaginal swabs are more sensitive than clinician taken endocervical culture for gonorrhoea
- e47** Compression-only CPR may improve survival for patients in cardiac arrest due to shockable rhythms treated by bystanders with public access defibrillation
- e48** Muscle strength during adolescence is associated with longevity
- e49** Child poisonings are more common in households experiencing psychosocial stressors
- e50** Use of psychotropic medication in Alzheimer's disease is associated with more rapid cognitive and functional decline

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator network

recycle

When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

