

Evidence-Based Medicine



Contents

Perspective

- 1 EBM, CME and the EMR
- 4 Loss of population data sources when health systems are not responsible for geographically defined populations: implications of the Health and Social Care Act of 2012 in England

Methods

- 6 Implementing GRADE: calculating the risk difference from the baseline risk and the relative risk

Therapeutics

- 9 Oral antibiotics confer small benefits and small harms in low-risk children with acute otitis media
- 10 A Mediterranean diet supplemented with olive oil or nuts reduces the incidence of major cardiovascular events in high-risk patients
- 11 Impact of honey as a topical treatment for wounds remains unclear
- 12 Medical management strategies to prevent recurrent nephrolithiasis are stagnant and stronger evidence is needed to reduce morbidity
- 13 Presence during cardiopulmonary resuscitation is beneficial to family members in the out-of-hospital setting
- 14 Normal diagnostic test results do not reassure patients
- 15 Combination therapy of steroids and antivirals improves the recovery rate in patients with severe Bell's palsy
- 16 Ramipril improves walking times and quality of life in patients with stable intermittent claudication
- 17 Prophylactic antibiotics for caesarean section administered preoperatively rather than post cord clamping significantly reduces the rate of endometritis
- 18 Limited impact of omega-3 fatty acids in patients with multiple cardiovascular risk factors
- 19 Self-management support has potential but is difficult to deliver in routine primary care
- 20 Lower target oxygen saturation levels in preterm infants are associated with increased mortality and decreased rates of retinopathy of prematurity
- 21 Conclusions about the benefits and risks associated with iron supplementation for anaemia prevention and control for preschoolers are limited by poor study quality
- 22 Dietary salt reduction; further lowering of target lowers blood pressure but may increase risk
- 23 Varenicline may reduce negative effect while aiding smoking cessation

Prognosis

- 24 Two-thirds of female smokers aged over 50 die from smoking-related causes, but quitting before 40 decreases this by 90%

February 2014 Volume 19 No 1

- 25 Weight gain does not attenuate cardiovascular benefits of smoking cessation

Aetiology


- 26 Perioperative β -blockade improves outcomes in higher risk patients following non-cardiac surgery
- 27 Maternal alcohol use increases risk of infant mortality
- 28 High-potency statins are associated with increased hospitalisations with acute kidney injury
- 29 Randomised controlled trial: extended-duration dabigatran is noninferior to warfarin and more effective than placebo for symptomatic VTE
- 30 Caffeine is modestly associated with lower birth weight and fetal growth
- 31 For proton pump inhibitor-dependent gastro-oesophageal reflux, laparoscopic fundoplication is superior to medical therapy at 5 years of follow-up
- 32 Need for standardising adverse event reporting in testosterone trials
- 33 Lower sodium intake reduces blood pressure in adults and children, but is not associated with a reduced risk of all CVD or all cause mortality
- 34 Sublingual immunotherapy improves symptoms of allergic rhinoconjunctivitis and asthma
- 35 Text messaging interventions increase adherence to antiretroviral therapy and smoking cessation
- 36 Increased cancer risk associated with CT in childhood
- 37 Use of narcotic analgesics associated with increased falls and fractures in elderly patients with osteoarthritis
- 38 Some more evidence of long-term psychosocial harms from receiving false-positive screening mammography results


Economic analysis

- 40 Conservative management for low-risk prostate cancer improves quality-adjusted life expectancy at lower cost compared with initial treatment

Online articles

- e1 Routine antibiotics given for uncomplicated severe acute malnutrition reduce mortality and improve nutritional recovery
- e2 Inconclusive evidence about the efficacy of diverse strategies for intermittent versus daily inhaled corticosteroids for persistent asthma in adults and children
- e3 Atrial fibrillation is associated with an increased risk of cognitive impairment and dementia with or without a history of clinical stroke
- e4 Maternal vitamin D insufficiency is associated with adverse pregnancy and neonatal outcomes
- e5 Short sleep duration is associated with increased BMI in adolescents over time
- e6 Long-term aspirin use and neovascular age-related macular degeneration: association or causation?

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

