

Evidence-Based Medicine

Contents

<p>Purpose and Procedure 2</p> <p>Additional Articles Abstracted in ACP Journal Club 3</p> <p>EBM NOTEBOOK</p> <p>Transferring evidence from research into practice: 2. Getting the evidence straight 4</p> <p>Announcing the arrival of Best Evidence 6</p> <p>Workshops for teaching evidence-based practice 7</p> <p>THERAPEUTICS</p> <p>Antibiotics were ineffective for the common cold unless bacteria were present in nasopharyngeal secretions 9</p> <p>Intranasal lidocaine relieved acute migraine pain 10</p> <p>Extracorporeal membrane oxygenation improved survival in mature newborn infants with severe respiratory failure 11</p> <p>Meta-analysis: Failure of oral rehydration therapy is infrequent in young, well-nourished children with gastroenteritis 12</p> <p>Imipramine was more effective than mianserin for nocturnal enuresis 13</p> <p>Meta-analysis: Intensive chemotherapy improves survival in childhood acute lymphoblastic leukaemia 14</p> <p>Review: Routine antimalarial chemoprophylaxis during pregnancy reduces parasitaemia and raises packed cell volume 15</p> <p>Terazosin reduced BPH symptoms and finasteride did not 16</p> <p>Combination therapy was better than zidovudine alone for HIV infection 17</p> <p>Streptokinase increased mortality in acute ischemic stroke 18</p> <p>Adjusted-dose warfarin for atrial fibrillation reduced stroke risk better than low-dose warfarin plus aspirin 19</p> <p>Extending enoxaparin 1 month after hospital discharge reduced thromboembolism after elective hip surgery 20</p>	<p>Bypass surgery and angioplasty led to similar 5-year mortality rates in multivessel coronary artery disease 21</p> <p>Meta-analysis: Antidyslipidemic therapy prevents myocardial infarction and death 22</p> <p>Meta-analysis: Dietary strategies alone reduce weight best in NIDDM 23</p> <p>Meta-analysis: <i>Hypericum</i> extracts (St. Johnswort) are effective for depression 24</p> <p>DIAGNOSIS</p> <p>Mini-Mental State Examination and the Informant Questionnaire on Cognitive Decline were efficient screening tests for dementia 25</p> <p>PROGNOSIS</p> <p>Children with perinatal HIV infection lived to an estimated mean age of 9.4 years 26</p> <p>ETIOLOGY</p> <p>Ruptured membranes, low CD4⁺ counts, low birthweight, and hard-drug use increased the risk for perinatal transmission of HIV-1 27</p> <p>Postmenopausal estrogen reduced the risk for and delayed the onset of Alzheimer disease 28</p> <p>Calcium antagonists did not increase mortality in patients with coronary artery disease 29</p> <p>ECONOMICS</p> <p>Health checks conducted by nurses in primary care cost between £1.46 and £2.25 per patient per 1% reduction in coronary risk 30</p> <p>A cardiovascular programme led by nurses cost between £5.08 and £5.78 per patient per 1% reduction in coronary risk 31</p> <p>Antenatal care by a general practitioner or midwife was less expensive than obstetrician-led shared care 32</p> <p>Glossary Inside back cover</p>
---	---