

Evidence-Based Medicine

Contents

Purpose and Procedure	34
Additional Articles Abstracted in ACP Journal Club	35
EBM NOTEBOOK	
Transferring evidence from research into practice: 3. Developing evidence-based clinical policy	36
... so little time, and	39
Corrections	40
THERAPEUTICS	
Co-amoxiclav had short-term effectiveness for persistent otitis media with effusion in young children ...	41
Review: Antibiotics are ineffective for acute bronchitis ...	42
Valaciclovir accelerated healing in recurrent herpes genitalis	43
Standardised pharmacotherapy and psychotherapy were more effective than usual care for major depression in primary care	44
Multifaceted intervention improved the treatment of depression in adults in a primary care setting	45
Emotionally focused marital therapy decreased marital distress and improved marital functioning in couples with ill children	46
Both sertraline and imipramine reduced symptoms in dysthymia	47
Meta-analysis: Training health professionals to provide smoking cessation interventions only slightly increases smoking cessation	48
Patients may overestimate their success in lowering coronary risk factors	49
Meta-analysis: Mortality is reduced when fibrinolytic therapy is started soon after the onset of MI symptoms ...	50
Lowering LDL cholesterol levels reduced fatal coronary events in patients with acute MI and average cholesterol levels	51
Meta-analysis: Thrombolytic therapy increases the risk for early death and intracranial hemorrhage after acute ischemic stroke	52
Endoscopic biliary endoprosthesis had a higher long-term complication rate than did duct clearance	53
Blunt-tipped needles reduced the incidence of glove puncture during abdominal closure	54
Hormone replacement therapy did not reduce the number of episodes of urinary incontinence	55
Fewer antenatal visits reduced high-risk referrals and preterm births in Zimbabwe	56
Meta-analysis: Misoprostol reduces NSAID-induced gastrointestinal mucosal injury	57
Meta-analysis: Respiratory rehabilitation relieves dyspnea in COPD	58
DIAGNOSIS	
D-dimer levels detected DVT in patients hospitalized for stroke rehabilitation	59
PROGNOSIS	
Ischemic stroke with accompanying atrial fibrillation was associated with reduced survival and functional status ...	60
ETIOLOGY	
Low cholesterol levels were associated with suicide in men	61
ECONOMICS	
Intensive therapy extended life and was cost-effective for IDDM	62
Cost-effectiveness of pravastatin in the secondary prevention of CAD in men varied with the risk profile of the patient	63
A day hospital and crisis respite program was less expensive than a hospital inpatient program in acute psychiatric illness	64
Glossary	Inside back cover