

Evidence-Based Medicine

Contents

Contents	129
Purpose and Procedure	130
Additional Articles Abstracted in <i>ACP Journal Club</i>	131
EBM NOTEBOOK	
Evidence-based pathology	132
Evidence-based medicine in primary care: some views from the 3rd U.K. workshop on teaching evidence-based medicine	133
Letter	135
THERAPEUTICS	
Nebulised ipratropium combined with albuterol led to improved pulmonary function in children with severe asthma	137
Intravenous salbutamol decreased recovery time in acute severe asthma in children	138
A nurse-led education programme reduced readmissions in children hospitalised with acute asthma	139
Review: Low-dose diuretics, but not high-dose diuretics and β -blockers, reduce coronary heart disease and total mortality	140
Digoxin reduced hospitalizations in patients with heart failure and normal sinus rhythm	141
Amiodarone reduced ventricular fibrillation or arrhythmic death in MI with frequent ventricular premature depolarizations	142
Amiodarone reduced arrhythmic death but not cardiac death in patients with MI and depressed left ventricular function	143
Compression stockings reduced the occurrence of the post-thrombotic syndrome in proximal DVT	144
Weight-loss and sodium-reduction programmes reduced BP and the risk for hypertension in overweight adults with high-normal BP	145
Quality-of-life measures improved in a mild hypertension study	146
Review: Laxatives and fibre therapies improve bowel movement frequency in adults with chronic constipation	147
A "critical time" intervention reduced homelessness in inner-city men	148
Review: Discontinuation rates are the same for serotonin-specific reuptake inhibitors and newer tricyclic and heterocyclic antidepressants	149
Review: School-based sex education linked to access to contraceptive services may reduce teenage pregnancy ...	150
Plasma exchange and intravenous immunoglobulin were equally effective for the Guillain-Barré syndrome	151
Amoxicillin did not improve the clinical course of acute maxillary sinusitis in primary care	152
Nitric oxide reduced the use of extracorporeal membrane oxygenation but not death in infants with hypoxic respiratory failure	153
DIAGNOSIS	
Ultrasonographic attenuation predicted fractures in older women	154
PROGNOSIS	
Residual leukaemia was not associated with relapse and was present during remission in children with acute lymphoblastic leukaemia	155
Review: Risk for suicide is increased for most mental disorders where patients require treatment in a hospital setting	156
ETIOLOGY	
Smoking increased the risk for death and MI after percutaneous revascularization	157
Review: Evidence relating sleep apnea to adverse health outcomes is inconclusive	158
Driving cessation was associated with depressive symptoms among older persons	159
ECONOMICS	
Additional cost per life-year saved with community thrombolysis, relative to hospital thrombolysis, was as low as £3890	160
Glossary	Inside back cover