

Evidence-Based Medicine



Contents

Perspectives

- 1 Peer review of human studies run amok: a break in the fiduciary relation between scientists and the public
 OPEN ACCESS
- 2 Harms of CT scanning prior to surgery for suspected appendicitis
 EDITOR'S CHOICE
- 3 Harms of CT scanning prior to surgery for suspected appendicitis
 EDITOR'S CHOICE

Therapeutics/Prevention

- 5 Naloxegol increases frequency of bowel movements and combats inadequate response to laxatives
- 6 ABT-450/r-ombitasvir and dasabuvir with ribavirin eliminates viraemia in most patients with HCV infection with cirrhosis
- 7 Zanamivir for influenza in adults and children shows limited benefit for treatment of symptomatic influenza and no effect on relevant complications
- 8 Effects of screening and lifestyle counselling on incidence of ischaemic heart disease in general population are limited
- 9 Phase 3 trial with the RTS,S/AS01 malaria vaccine shows protection against clinical and severe malaria in infants and children in Africa
- 10 Antithrombotic therapy for pregnancy complications: let's not throw the baby out with the bath water
- 11 Updated review identifies no adverse impact on mother or offspring during the perinatal period of aspirin use for prevention of preeclampsia
- 12 With statin co-administration, drugs designed to increase HDL have no impact on cardiovascular outcomes
- 13 Calcaneal fractures have universally poor outcomes regardless of management
- 14 Fruit and vegetable consumption is associated with reduced all-cause and cardiovascular mortality
- 15 Delaying initiation of ART for 5 weeks improves survival in patients with HIV infection with cryptococcal meningitis
- 16 Epidural steroid injections are not effective for patients with lumbar spinal stenosis
- 17 Initial choice of medication has little effect on short-term or long-term outcome for most patients with Parkinson's disease
- 18 Gastric bypass has better long-term outcomes than gastric banding
- 19 Pulse oximetry may lead to unnecessary hospital admissions for infants with bronchiolitis and mild hypoxaemia
- 20 Collaborative care improves clinical outcomes for adolescents with depression treated in primary care
- 21 Blood pressure-lowering treatment should be based on the level of cardiovascular risk, not on the level of blood pressure
- 22 Gum chewing is associated with early recovery of bowel motility and shorter length of hospital stay for women after caesarean section
- 23 Treatment of hepatitis C, then, now and tomorrow

February 2015 Volume 20 No 1

- 24 The therapeutic value of hypertonic saline in acute bronchiolitis remains unclear

Prognosis

- 26 For patients who underwent elective non-cardiac surgery, a history of stroke is associated with an increased risk of major adverse cardiovascular events and death, particularly if time elapsed between stroke and surgery is less than 9 months
- 27 Aiming too high or too low? Searching for the appropriate therapeutic thresholds in hypertension is not over yet
- 28 The unprecedented scale of the West African Ebola virus disease outbreak is due to environmental and sociological factors, not special attributes of the currently circulating strain of the virus
- 29 Long-term follow-up after removal of colorectal adenomas provides evidence for risk stratification of patients at colonoscopic polypectomy
- 30 Study results suggest a positive effect of light-to-moderate physical activity on disability among those with or at risk of knee arthritis

Diagnosis

- 31 Evidence is lacking to support pelvic examinations as a screening tool for non-cervical cancers or other conditions
- 32 CT should not be relied on for cases of isolated vomiting in children with blunt head trauma
- 33 Physician practice and PECARN rule outperform CATCH and CHALICE rules based on the detection of traumatic brain injury as defined by PECARN

Aetiology/Harm

- 35 Pertussis: a significant cause of prolonged acute cough in previously vaccinated school-aged children
- 36 Risk of warfarin-associated intracerebral haemorrhage after ischaemic stroke is low and unchanged during the 2000s
- 37 Potential confounding still clouds the possible association of maternal caffeine intake and low birth weight
- 38 Mendelian randomisation meta-analysis sheds doubt on protective associations between 'moderate' alcohol consumption and coronary heart disease
 EDITOR'S CHOICE

Letters

- 39 Low-dose opioids should be considered for symptom relief also in advanced chronic obstructive pulmonary disease (COPD)
- 40 Response to Ekström *et al* regarding low-dose opioids in advanced chronic obstructive pulmonary disease

Corrections

- 34 Corrections

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator network

recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

