

Evidence-Based Medicine



Contents

June 2015 Volume 20 No 3

Original EBM research

- 81** Impact of a GRADE-based medical question answering system on physician behaviour: a randomised controlled trial

Systematic review

- 88** Special tests for assessing meniscal tears within the knee: a systematic review and meta-analysis

Therapeutics/Prevention

- 98** There is no good evidence for the effectiveness of commonly used over-the-counter medicine to alleviate acute cough
- 99** Prolonged dual antiplatelet therapy after drug-eluting stent reduces the risk of stent thrombosis and major cardiovascular events but increases the rate of bleeding
- 100** Up to 4000 mg of paracetamol a day is ineffective for acute low back pain
- 101** Using the Watchman device to close the left atrial appendage reduces risk of stroke in atrial fibrillation, compared to using warfarin
- 102** Although the evidence is not clear, decreases in prostate cancer mortality in specific subgroups of men may be due to screening
- 103** Weight-loss diets only work when you follow them
- 105** Postfracture pain in children can be adequately managed with ibuprofen
- 106** Nebulised hypertonic saline does not reduce hospital length of stay in acute bronchiolitis
- 107** Coronary artery bypass graft surgery provides clinically important quality-of-life improvements over medical therapy in patients with ischaemic left ventricular dysfunction
- 108** Timing of thrombolysis for acute ischaemic stroke: the earlier the treatment the better the outcome, irrespective of age or stroke severity
- 109** Invasive revascularisation in patients with moderate intermittent claudication provides a significant improvement in quality of life compared with conservative treatment
- 110** Monthly malaria chemoprevention shows potential in an area of very high, perennial malaria transmission

Prognosis

- 111** Angina may trump ischaemia in predicting stable coronary artery disease outcomes, but most cardiovascular events occur in patients without angina or asymptomatic ischaemia
- 112** Violent reinjury and mortality highlights the need for a comprehensive care approach to youth presenting for assault-related injury

Aetiology/Harm


- 113** Cotrimoxazole is associated with sudden death in older patients receiving inhibitors of renin-angiotensin system
- 114** Maternal and neonatal outcomes in nulliparous women undergoing non-medically indicated induction of labour or expectant management at 38–41 weeks gestation
- 115** Metformin should not be contraindicated in patients with type 2 diabetes and mild to moderate renal impairment
- 116** The human papillomavirus vaccination is not associated with risk of multiple sclerosis or other demyelinating diseases
- 117** Vitamin K antagonist-experienced patients with a history of stroke/transient ischaemic attack who switched from warfarin to dabigatran increased their rate of recurrent stroke/transient ischaemic attack compared with those on warfarin
- 118** Prenatal and postnatal exposure to parental smoking increases odds of allergic diseases during childhood and adolescence


Diagnosis

- 119** Digital rectal examination can detect early prostate cancer
- 120** Symptoms alone may lead to inaccurate diagnosis in men with possible bladder outlet obstruction

PostScript

- 104** Correction

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

