

# Evidence-Based Medicine



## Contents

### Editorial

- 191** Address persistent racial disparities in academic medicine to improve healthcare quality



### Perspective

- 193** Not a Humbug: the evolution of patient-centred medical decision-making



### Therapeutics/Prevention

- 198** Induction of labour at 37–38 weeks in women with large fetuses decreases the likelihood of shoulder dystocia; however, overall benefit of early-term delivery has not been demonstrated

- 199** Low-molecular-weight heparin offers no benefit to patients with unexplained recurrent miscarriage in future pregnancy

- 200** Interruption of all anticoagulation is non-inferior to the use of short-term parenteral bridging in patients with atrial fibrillation undergoing invasive procedures

- 201** US counties with higher rates of breast cancer screening have higher rates of incidence with no concomitant decrease in breast cancer mortality suggesting overdiagnosis

- 202** Mediterranean diet may reduce Alzheimer's risk

- 203** Subcutaneous liraglutide reduces weight and improves metabolic control in obese participants

- 204** Early introduction of peanut to infants at high allergic risk can reduce peanut allergy at age 5 years

- 205** Paracetamol is ineffective for spinal pain and knee and hip osteoarthritis

- 206** Simplified sleep restriction impacts objective but not subjective sleep for people with primary insomnia in primary care

- 207** Synbiotic yogurt consumption leads to statistical improvements in child health that may not be clinically relevant

- 208** Routine iron supplementation for non-anaemic pregnant women in developed countries needs evaluation

- 209** Intra-arterial thrombectomy improves functional outcome when administered up to 6 h after stroke

- 210** Vaginal hysterectomy is the best minimal access method for hysterectomy

- 211** The success of mechanical thrombectomy in acute ischaemic stroke is strictly dependent on ischaemic core size and time to treatment

- 212** At 5 years, transcatheter aortic valve replacement had similar rates of mortality and stroke as surgical aortic valve replacement in high-risk patients

## December 2015 Volume 20 No 6

- 214** Meta-analysis based on limited data shows no evidence to support the guideline recommendation for early administration of antibiotics in severe sepsis and septic shock

- 215** A mandibular advancement device did not affect daytime sleepiness and quality of life in obstructive sleep apnoea

### Prognosis

- 217** Primary prevention implantable cardioverter defibrillators are effective in preventing overall mortality in older patients

- 218** Prior stillbirth is a risk factor for stillbirth in a subsequent pregnancy

### Aetiology/Harm

- 219** Postnatal environmental tobacco smoke exposure is associated with objective markers of atopy in preschool-aged children

- 220** Risks of transvaginal mesh may be associated with an increased risk of urinary retention and reintervention

- 221** Intermediate and long-term cognitive effects in older adults secondary to cardiovascular procedures is uncommon but current evidence is insufficient

- 222** Risk of new-onset diabetes with statin use should not be overemphasised

- 223** Sugar-sweetened beverage consumption is linked to global adult morbidity and mortality through diabetes mellitus, cardiovascular disease and adiposity-related cancers

### Diagnosis

- 225** Clinical outcomes following coronary CT angiography are comparable to radionuclide myocardial perfusion imaging for ethnically diverse intermediate risk acute chest pain inpatients

### Practice guideline

- 227** Guidelines for the management of extremely preterm deliveries in the grey zone of viability between 23 and 24 weeks' gestation vary widely in developed countries

### Economic analysis

- 228** Delayed hospital admission until active labour may help prevent unnecessary caesarean delivery

- 229** Paediatric pain-related conditions impact healthcare expenditures

- 230** Current 10-year atherosclerotic cardiovascular disease risk threshold for statin eligibility is cost-effective for primary prevention

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org.uk](http://www.publicationethics.org.uk)

equator network

recycle  
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

