


Evidence-Based Medicine



Contents

Perspective

- 201 'Spin' in reports of clinical research



Original EBM Research

- 203 Avoiding alert fatigue in pulmonary embolism decision support: a new method to examine 'trigger rates'

Methods

- 208 Value and usability of unpublished data sources for systematic reviews and network meta-analyses



Systematic review

- 214 Non-pharmacological treatment of depression: a systematic review and evidence map

Therapeutics/Prevention

- 222 LABA/LAMA combinations instead of LABA/ICS combinations may prevent or delay exacerbations of COPD in some patients
- 223 Most add-on therapies to metformin have similar effects on HbA1c
- 224 Existing evidence is insufficient to justify metformin or other agents as first-line therapy for type 2 diabetes
- 225 Aspirin reduces cardiovascular events in primary prevention of cardiovascular disease but at a near equivalent risk of increased bleeding
- 226 Uncertainty in evidence synthesis limits clinical applicability of a clinical and cost-effectiveness analysis of induction of labour methods
- 227 Compared to conventional CPR for in-hospital cardiac arrest, extracorporeal-CPR is associated with improved survival to hospital discharge and more favourable neurological outcome
- 228 In individuals at intermediate risk for cardiovascular disease, treatment with rosuvastatin but not candesartan plus hydrochlorothiazide lowers cardiovascular disease event rates

December 2016 Volume 21 No 6

Prognosis

- 230 Rate of epilepsy in people with autism and the rate of autism in people with epilepsy are high
- 231 Implementing evidence-based practices improves neonatal outcomes

Aetiology/Harm

- 232 Low-dose second-generation oral contraceptives are associated with the lowest increased risk of cardiovascular adverse effects
- 233 Careful monitoring of fetal growth and maternal nutritional status should be practiced in pregnant women with a history of bariatric surgery
- 234 Morphine exposure in preterm infants correlates with impaired cerebellar growth and poorer neurodevelopmental outcome
- 235 Cognitive adverse effects and brain deterioration associated with use of anticholinergic activity medicines in older adults

Diagnosis

- 236 Traditionally taught clinical variables and risk factors perform poorly in the prediction of acute coronary syndromes in the emergency department

Practice guideline

- 237 Determination of a single, universal threshold for caesarean section rate is not the way forward



Letters

- 238 Evidence pyramids
- 238 Letter in reply to 'Evidence pyramids' from Dr Kaufmann
- 239 Letter in reply to 'Evidence pyramids' from Dr Kaufmann
- 240 Where should preappraised evidence summaries and guidelines place in a pyramid?



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
Member since 2010
JM0659
www.publicationethics.org.uk

equator
network

recycle
When you have finished
with this please recycle it

Receive regular table of contents by email.
Register using this QR code.

