

## EBM analysis

# Transparency of COVID-19 vaccine trials: decisions without data

Sarah Tanveer <sup>1</sup>, Anisa Rowhani-Farid <sup>1</sup>,  
Kyungwan Hong,<sup>1</sup> Tom Jefferson <sup>2</sup>, Peter Doshi <sup>1</sup>

10.1136/bmjebm-2021-111735

<sup>1</sup>Department of Pharmaceutical Health Services Research, University of Maryland Baltimore, Baltimore, Maryland, USA  
<sup>2</sup>Department for Continuing Education, University of Oxford, Oxford, UK

Correspondence to:  
**Dr Peter Doshi**,  
Pharmaceutical Health Services Research, University of Maryland School of Pharmacy, Baltimore, MD 21201, USA; pdoshi@rx.umaryland.edu

## Transparency in clinical trials: an established norm across sectors

Access to data for drugs and vaccines has historically been fairly limited to journal article publications and hard-to-access and difficult to read regulatory reports.<sup>1</sup> But the past decade has witnessed strides in clinical trial data transparency. A wide range of institutions, from pharmaceutical companies, government agencies, trade organisations, journals and not-for-profit organisations, have all acknowledged the importance of data sharing, including the release of deidentified individual participant data. Many policies, regulations and platforms now exist to facilitate data access, including landmark transparency policies from the European Medicines Agency (EMA)<sup>2,3</sup> and Health Canada.<sup>4</sup> Both regulators now post on their websites, sections of the licensure dossier received by the industry (<https://clinicaldata.ema.europa.eu/> and <https://clinical-information.canada.ca/>). There are also industry and academic platforms to facilitate third-party access to trial data and documents, including ClinicalStudyDataRequest.com, Yale University Open Data Access (YODA) Project and Vivli.<sup>5</sup> In 2013, the US and European industry trade organisations endorsed a joint statement on clinical trial data sharing, making a series of commitments that ‘recognise the importance of sharing clinical trial data in the interest of patients, healthcare and the economy’<sup>6</sup> In 2015, the US Institute of Medicine similarly endorsed benefits of sharing clinical trial data, emphasising that ‘verification and replication of investigators’ claims’ were essential to the scientific process, and noting the numerous benefits to stakeholders ‘including payers of healthcare as well as patients, their physicians and researchers.’<sup>7</sup>

## Why we need access to COVID-19 vaccine trial data and documents

Clinical trial transparency is always important, but is especially critical during the COVID-19 pandemic (or any public health emergency) where regulatory decisions are being made quickly by government health officials, novel vaccine platforms are being used, vaccines are being administered widely and taxpayer funds have contributed heavily to research and development. Critical appraisal of clinical trials is vital to inform decision making at the personal, professional

## Summary box

- ⇒ Data transparency has become a well-established norm in biomedical research, and is especially important for broadly used public health interventions like COVID-19 vaccines.
- ⇒ Tax payers helped fund COVID-19 vaccine trials and should have the right to access the results.
- ⇒ There is inadequate availability of COVID-19 vaccine trial documents and data; individual participant data will not be available for months, perhaps years, for most vaccines.
- ⇒ Widespread use of interventions without full data transparency raises concerns over the rational use of COVID-19 vaccines.
- ⇒ Trial transparency must start early and be continuous. Trial protocols should be released once finalised, before trial results are reported, and should be accompanied with the release of trial documents and data before clinicians and the public make decisions regarding product use.

and governmental level, but cannot be credibly performed on journal publications alone.<sup>8,9</sup> Access to clinical trial data and related trial documents (see [box 1](#)) allows for independent and informed assessment of trials. By understanding details of how studies were designed and how data were collected, one can understand if endpoints were reliably operationalised and measured. Similarly, release of underlying data from clinical trials allows for independent verification of results, assessment of heterogeneity of treatment effects for specific subgroups, and facilitates the formation of new research questions.<sup>10</sup>

There are specific issues in COVID-19 vaccine trials that merit scrutiny. Consider blinding, an essential feature in randomised trials investigating efficacy against subjective endpoints, as in the COVID-19 vaccine trials. Assessing the reliability of blinding involves analysing endpoint definitions and data collection. The primary endpoint in many trials is laboratory-confirmed, symptomatic



© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Tanveer S, Rowhani-Farid A, Hong K, *et al.* *BMJ Evidence-Based Medicine* 2022;**27**:199–205.

## Box 1 Types of trial documents

**Case report forms (CRFs)**

The original paper or electronic forms on which individual participants' data (demographic, efficacy measurements, adverse events, etc) are recorded during the clinical trial. These documents contain structured fields which make it easier to analyse and report trial data. Access to blank case report forms (CRFs) allows for independent evaluation of how data were collected and endpoints were operationalised.

**Clinical study reports (CSRs)**

Unabridged, structured report of a clinical study written for regulators.<sup>30</sup> A complete clinical study reports (CSRs), including study appendices, includes documentation of trial design, results of trial included adverse events, trial protocol, statistical analysis plans and blank CRFs. CSRs on average span thousands of pages, making them a rich source of information. European Medicines Agency and Health Canada are the only two regulatory agencies that publish CSRs at this time.

**Certificate of analysis**

Provides a description of the chemical analysis and physical appearance of the study interventions actually used in the trial (both experimental and comparators).

**Protocol**

Document written prior to study start date which details plans on how the study will be conducted, analysed and reported. Any changes or deviations from the trial protocol should be tracked and provided, with a rationale for the change, in the form of a formal protocol amendment.

**Statistical analysis plan (SAP)**

A written plan of how trial data will be analysed and which statistical methods and definitions will be used.

**Informed consent form (ICF)**

A document required to be provided to study subjects that contains information related to the description of the study, purpose, study intervention(s), any procedures, adverse events, risk and benefits, compensation and rights of participants enrolled in the study.

**Serious adverse event (SAE) narratives**

Unstructured paragraphs of text providing details and context of the serious adverse events that occurred in study participants. Narratives are usually contained within a CSR.

**Electronic individual participant data (IPD)**

Complete electronic computerised dataset for each participant in the trial which allows for full replication of study findings using statistical software. Complete CSRs also contain participant level data, mostly in appendices, but these are in text (not dataset) form as individual line listings.

**Investigational Medicinal Product Dossier (IMPD)**

Continued

## Box 1 Continued

Document that describes the quality of the placebo and investigational product, how the product(s) were manufactured, non-clinical and clinical study results. An Investigational Medicinal Product Dossier is required for all clinical trials conducted in the European Union.

**Investigator's Brochure (IB):**

A living document containing a summary of the clinical and nonclinical data of an investigational product, including its pharmacology, pharmacokinetics, toxicology and adverse event profile, among other items.

Sources: Restoring Invisible and Abandoned Trials (RIAT) declaration,<sup>31</sup> RIAT Support Center Glossary.<sup>32</sup>

COVID-19. However, prior to the release of trial protocols, few details were known about this endpoint. Registry entries were vague (eg, Pfizer's largest study<sup>11</sup> only stated 'confirmed COVID-19' as one of 35 primary outcome measures), leaving unclear how the definition was operationalised. While the subsequent release of some protocols addressed some questions, it raised new questions that can only be answered with underlying data. Protocols make clear that the symptomatic component of the primary endpoint was reported by trial participants, typically via a smartphone app, and defined by one or more signs and symptoms, many of which were subjective (eg, in Pfizer's trial, COVID-19 symptoms included at least one of the following: fever, cough, shortness of breath, chills, muscle pain, loss of taste or smell, sore throat, diarrhoea or vomiting.) The fact that the placebo was saline and the vaccines cause short term adverse events in the majority of people raises concerns about unofficial unblinding—that is, the ability of trial participants and investigators to make educated guesses as to treatment allocation. Only a thorough analysis of the underlying individual participant data will allow for an exploration of the extent to which unofficial unblinding may have occurred and biased data collection for the primary endpoint. Access to data would also allow for straightforward replication studies, perhaps particularly important when real-world results appear incompatible with reported trial results. For example, at the time of writing (27 June 2021), Seychelles, Mongolia, Bahrain, Uruguay, and Chile were experiencing COVID-19 outbreaks despite high uptake of WHO-authorised vaccines.

In addition, trial protocols from Pfizer and Moderna indicate that event adjudication committees were involved in counting COVID-19 cases. Considering that the primary endpoint was defined as a positive lab test and patient-reported symptoms, it is unclear how an adjudication committee might affect the primary endpoint evaluation process. Transparency of the committee's charter may provide additional detail on what data committee members had access to in forming their judgements (eg, did they have access to data on patients' symptoms in the first week after vaccination, when vaccine-related adverse events could be expected?), and what criteria they used to form their judgements. Access to such documents, therefore, is also important.

Regarding adverse events, detailed narratives of serious adverse events that occurred in a trial are a standard element found within clinical study reports and can enable a more thorough understanding of potential harms. Patterns in adverse

**Table 1** Currently available COVID-19 vaccine trial data for selected trials

Trial ID; no enrolled; included ages	Pre-study documents	Post-study documents†					Total pages available§
		Press release	Pub	CSR	Other‡	IPD	
<b>Pfizer BNT162b2 mRNA vaccine</b>							
<a href="#">NCT04368728</a> ; n=43 998; 12–85 years	Protocol, SAP, Blank CRF	Press release 1, 2, 3	Pub 1,2,3	CSR	Other	No	3880
<a href="#">NCT04713553</a> ; n=1530; 12–50 years	None	No	N/A: trial ongoing				0
<a href="#">NCT04816643</a> ; n=4644; 6 months to 11 years	None	No	N/A: trial ongoing				0
<b>Moderna mRNA-127 vaccine</b>							
<a href="#">NCT04470427</a> ; n=30 420; ≥18 years	Protocol, SAP	Press release	Pub	No	Other	No	3293
<a href="#">NCT04811664</a> ; n=37 500; 18–26 years	None	No	N/A: trial ongoing				0
<a href="#">NCT04796896</a> ; n=6750; 6 months to 12 years	None	No	N/A: trial ongoing				0
<b>Oxford/AstraZeneca ChAdOx1 vaccine</b>							
<a href="#">ISRCTN89951424</a> ; n=10 300; ≥18 years	Protocol	Press release	Pub 1¶, 2¶	No	No	No	123
<a href="#">NCT04400838</a> ; n=12 390; ≥18 years	Protocol	Press release	Pub 1¶, 2¶	No	No	No	214
<a href="#">ISRCTN15638344</a> ; n=300; 6–17 years	None	No	N/A: trial ongoing				0
<b>Janssen (Johnson &amp; Johnson) Ad26.COVS vaccine</b>							
<a href="#">NCT04505722</a> ; n=44 325; ≥18 years	Protocol, SAP, Blank ICF	Press release	Pub	No	No	No	530
<a href="#">NCT04535453</a> ; n=1210; 12 to 55, ≥65 years	None	No	N/A: trial ongoing				0
<a href="#">NCT04614948</a> ; n=30 000; ≥18 years	Protocol	No	N/A: trial ongoing				166
<b>Novavax SARS-CoV-2 rS/Matrix-M1 Adjuvanted vaccine</b>							
<a href="#">NCT04611802</a> ; n=30 000; ≥18 years	Protocol	No	N/A: trial ongoing				128
<a href="#">NCT04368988</a> ; n=1419; 18–84 years	Protocol, SAP	Press release	N/A: trial ongoing				189
<a href="#">NCT04583995</a> ; n=15 187; 18–84 years	Protocol, SAP	Press release	Pub	No	No	No	128
<b>Gamaleya Research Institute Sputnik V/Gam-COVID-Vac vaccine</b>							
<a href="#">NCT04530396</a> ; n=33 758; ≥18 years	None	Press release	Pub	No	No	No	11
<a href="#">NCT04741061</a> ; n=6000; ≥18 years	None	No	N/A: trial ongoing				0
<a href="#">NCT04642339</a> ; n=2000; ≥18 years	None	No	N/A: trial ongoing				0
<b>Sinopharm (BIBP) vaccine</b>							
<a href="#">ChiCTR2000032459</a> ; n=2128; ≥3 years	None	No	Pub	No	No	No	13
<a href="#">NCT04510207</a> ; n=45 000; ≥18 years	Protocol, SAP	No	Pub	No	No	No	102
<a href="#">NCT04612972</a> ; n=12 000; ≥18 years	None	No	N/A: trial ongoing				0
<b>Sinovac (CoronaVac) vaccine</b>							
<a href="#">NCT04456595</a> ; n=12 688; ≥18 years	Protocol	Press release	Pub	No	No	No	201
<a href="#">NCT04551547</a> ; n=552; 3–17 years	None	No	Pub	No	No	No	22
<a href="#">NCT04582344</a> ; n=13 000; 18–59 years	Protocol	Press release	No	No	No	No	57

Data current as of 27 June 2021.

\*Pre-study documents include: protocol, statistical analysis plan, blank informed consent form, blank case report form, data monitoring board charter, event adjudication committee charter, investigational medicinal product dossier and investigator's brochure.

†Post-study documents include: press releases (that contain any results), journal publication (including pre-prints), clinical study report and individual participant data.

‡Other includes documents released by Health Canada and EMA other than the CSR.

§Total pages available excludes press releases. Access to the dataset used to determine page count for trials where additional data were available through Health Canada and the European Medicines Agency is available in the Zenodo repository (<http://doi.org/10.5281/zenodo.4737417>).

¶Pooled trial analysis publication listed if there were no individual trial publications.

CRF, case report form; CSR, clinical study report; EMA, European Medicines Agency; ICF, informed consent form; IPD, individual participant data; n, number enrolled in trial; N/A, not applicable; Pub, journal publication; SAP, statistical analysis plan.

events can be explored through access to electronic individual participant-level data.

#### What data have been released

For eight COVID-19 vaccines being used, or under consideration for use globally (Pfizer, Moderna, Oxford/AstraZeneca, Janssen/

Johnson & Johnson, Novavax, Gamaleya Institute, Sinopharm and Sinovac), we evaluated the public availability of a variety of important pre-study documents (eg, trial protocol, statistical analysis plan, blank informed consent form, blank case report form, data monitoring board charter, and event adjudication committee charter) and post-study documents (eg, press releases with trial

**Table 2** Timing of release of individual participant data from COVID-19 vaccine trials

Phase 3 trial	Protocol released before results released?	Pledge to share IPD	Estimated date of availability (based on data sharing statement in protocol or publication)
Pfizer phase 2/3; 43 998 participants (NCT04368728)	Yes	Yes	April 2025, based on statement in trial protocol that data will be made available '24 months after study completion'
Moderna phase 3; 30 420 participants (NCT04470427)	Yes	Unclear	October 2022, based on statement in trial publication that data "may be available ... once the trial is complete"
Oxford/AstraZeneca phase 3; 10 300 participants (ISRCTN89951424)	No	Yes	December 2021, based on statement in trial publication that trial data 'will be made available when the trials are complete'
Janssen (Johnson & Johnson) phase 3; 44 325 participants (NCT04505722)	Yes	Yes	Unclear. April 2021 publication suggested data availability will begin 'with publication,' but as of June, still not listed on Yale Open Data Access Project website.
Novavax phase 3; 30 000 participants (NCT04611802)	Yes	No*	We could not locate any other written statement regarding patient-level data sharing. It is not discussed in the trial protocol or publication.
Gamaleya Research Institute phase 3; 33 758 participants (NCT04530396)	No	Yes*	May 2021, based on statement in trial publication that data will be made available 'on completion of clinical trials'
Sinopharm phase 3; 45 000 participants (NCT04510207)	No	Yes	December 2022, based on statement in trial publication that data will be available between December 2022 and December 2027, with reasonable request to the sponsor and principal investigator.
Sinovac phase 3; 13 000 participants (NCT04582344)	Yes	No*	We could not locate any other written statement regarding patient-level data sharing. It is not discussed in the full-length study protocol. Also, a structured summary of study protocol states 'Not applicable' under the availability of data and material.

Data current as of 27 June 2021.  
\*According to the 'Plan to Share IPD' field in the ClinicalTrials.gov entry.  
IPD, individual participant data.

results, journal publication, clinical study report and individual participant data availability) at the time of writing (27 June 2021). We counted the total number of pages available as a crude proxy for the level of detail, as some documents, like clinical study reports, can be highly variable in length depending on the availability of appendices.

The overall picture is one of varied transparency. While several trials have at this point published protocols and statistical analysis plans along with the study publications, with some even released while the trials were underway, many key trial documents remain inaccessible (table 1). For example, a WHO report found that out of 86 clinical trials for 20 COVID-19 vaccines, 12% of clinical trial protocols were made publically available.<sup>12</sup> In our analysis, trial protocols and informed consent forms were not available for trials involving special populations such as children (NCT04816643) and pregnant women (NCT04754594). And despite vaccine roll-out, electronic individual participant data is not available for most trials.<sup>12</sup> Some sponsors, such as Moderna, have sent mixed messages on whether they even intend to share data, while others, such as Pfizer and Sinopharm, indicate they will not even begin accepting data requests for many months or years (table 2). In other cases, trialists have indicated very narrow time frames for sharing data, for example, 'beginning 3 months and ending 1 year after publication' (ChiCTR2000032459).<sup>13</sup>

The greatest availability of data at present comes from the EMA, Health Canada, and Japanese Pharmaceuticals and Medical Devices Agency which have released thousands of pages from company submissions for COVID-19 vaccines, far exceeding what is available elsewhere. The Food and Drug Administration (FDA) does not routinely make any industry documents it receives publicly available. It should be noted, however, that the EMA and Health Canada's commitment to transparency does not necessarily equate with the public availability of critical study documents.

Judging from the availability of documents, regulators themselves appear to be receiving less complete and granular data than normal due to the compressed timeline from study start to regulatory decision making. For example, in July 2020, Canada granted emergency approval for the use of remdesivir, a drug used for the treatment of COVID-19, without receiving a copy of the clinical study report.<sup>14</sup> This is also the case for the Moderna vaccine, for which a variety of trial documents have been posted by Health Canada and the EMA, but not the clinical study report.<sup>15</sup> For Cuba, Russia, India and China state-developed vaccines, access to data prior to regulatory approval is even less clear.<sup>16</sup>

**The world's most used COVID-19 vaccines: Sinovac and Sinopharm**  
Sinovac and Sinopharm, developed by Chinese pharmaceutical companies, account for the majority of vaccines being in Asia, South America, the Caribbean and Africa.<sup>17</sup> They are authorised by the WHO and included in the WHO COVID-19 Vaccines Global Access initiative. However, transparency of trial data and documents is extremely limited, similar to other COVID-19 vaccines. Because neither Chinese vaccine has thus far been authorised by the EMA, Health Canada, or the Japanese Pharmaceuticals and Medical Devices Agency (the three regulators publishing data to their websites), there is also no expectation that data on these vaccines will become publicly available via regulators (table 3). Sinovac vaccines have been administered for nearly 1 year (since July 2020), and still have yet to publish clinical trial data (as at 27 June 2021). Trial results have largely been limited to government media reports and press releases.<sup>18</sup>

#### Transparency of regulatory decision making

Apart from data and documents tied to a specific trial, credible analysis and interpretation of data may require understanding regulatory decision making. For example, the fact that many

**Table 3** Availability of regulatory reviews and additional data for COVID-19 vaccines

Manufacturer	Regulatory reviews and additional data
Pfizer	<b>USA FDA:</b> <a href="#">main webpage</a> , <a href="#">Emergency Use Authorisation review memorandum</a> (57 pages), <a href="#">EUA review memorandum for 12–15 year olds</a> (43 pages), <a href="#">advisory committee briefing documents (FDA, Pfizer)</a> <b>UK MHRA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (51 pages) <b>AUS TGA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (42 pages) <b>EU EMA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (140 pages), <a href="#">clinical information</a> (6990 pages) <b>Canada HC:</b> <a href="#">main webpage</a> , <a href="#">summary basis of decision</a> (63 pages), <a href="#">clinical information</a> (5910 pages) <b>Japan PMDA:</b> <a href="#">main webpage (English, Japanese)</a> , <a href="#">public assessment report</a> (74 pages), <a href="#">clinical and non-clinical information</a> (1239 pages) <b>WHO:</b> <a href="#">main webpage</a> , <a href="#">background document to the WHO interim recommendations</a> (44 pages)
Moderna	<b>USA FDA:</b> <a href="#">main webpage</a> , <a href="#">Emergency Use Authorisation review memorandum</a> (61 pages), <a href="#">advisory committee meeting briefing documents (FDA, Moderna)</a> <b>UK MHRA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (7 pages) <b>EU EMA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (169 pages), <a href="#">clinical information</a> (6095 pages) <b>Canada HC:</b> <a href="#">main webpage</a> , <a href="#">summary basis of decision</a> (61 pages), <a href="#">clinical information</a> (6095 pages) <b>Japan PMDA:</b> <a href="#">main webpage (English, Japanese)</a> , <a href="#">public assessment report</a> (74 pages), <a href="#">clinical and non-clinical information</a> (542 pages) <b>WHO:</b> <a href="#">main webpage</a> , <a href="#">background document to the WHO interim recommendations</a> (41 pages)
Oxford/AZ	<b>UK MHRA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (58 pages) <b>AUS TGA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (49 pages) <b>EU EMA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (181 pages) <b>Canada HC:</b> <a href="#">main webpage</a> , <a href="#">summary basis of decision</a> (66 pages) <b>Japan PMDA:</b> <a href="#">main webpage (English, Japanese)</a> , <a href="#">public assessment report</a> (122 pages), <a href="#">clinical and non-clinical information</a> (1134 pages) <b>WHO:</b> <a href="#">main webpage</a> , <a href="#">background document to the WHO interim recommendations</a> (56 pages)
Janssen	<b>USA FDA:</b> <a href="#">main webpage</a> , <a href="#">Emergency Use Authorisation review memorandum</a> (69 pages), <a href="#">advisory committee meeting briefing documents (FDA, Janssen)</a> <b>UK MHRA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (11 pages) <b>EU EMA:</b> <a href="#">main webpage</a> , <a href="#">public assessment report</a> (218 pages) <b>Canada HC:</b> <a href="#">main webpage</a> , <a href="#">summary basis of decision</a> (60 pages) <b>WHO:</b> <a href="#">main webpage</a> , <a href="#">background document to the WHO interim recommendations</a> (54 pages)
Novavax	None
Gamaleya Research Institute	None
Sinopharm	<b>WHO:</b> <a href="#">main webpage</a> , <a href="#">background document to the WHO interim recommendations</a> (23 pages)
Sinovac	<b>WHO:</b> <a href="#">main webpage</a> , <a href="#">background document to the WHO interim recommendations</a> (30 pages)

Data current as of 27 June 2021.  
Regulatory agencies for each country.: USA FDA; UK MHRA; Canada HC; Japan PDMA; International/non-country entity WHO.  
AUS, Australia; EMA, European Medicines Agency; EU, European Union; FDA, Food and Drug Administration; HC, Health Canada; MHRA, Medicines and Healthcare products Regulatory Agency; PMDA, Pharmaceuticals and Medical Devices Agency; TGA, Therapeutic Goods Administration; USA, United States of America; WHO, World Health Organization.

experts had originally believed the trials were designed to study a reduction in hospitalisation, intensive care utilisation and death,<sup>19</sup> points to a need to better understand the rationale for primary endpoint selection. Regulators played a major role in shaping this. An FDA guidance document from June 2020, before phase 3 trials commenced, stated that laboratory-confirmed COVID-19 (of essentially any severity) was an acceptable primary endpoint. But there remains limited transparency on the rationale for the selection of this endpoint: why was it chosen? What other endpoints were considered? More information about the internal deliberations is essential to understand whether the decisions made were reasonable.<sup>20</sup>

Additionally, there is a need for transparency around any deliberations regarding the length of follow-up necessary to adequately assess efficacy and safety prior to licensure. Longer follow-up for investigational products such as COVID-19 vaccines and therapeutic agents are necessary to evaluate duration of protection and ensure public and professional confidence.<sup>21</sup> The International Coalition of Medicines Regulatory Authorities (ICMRA), a global collaborative coalition of 30 medicine regulators including the FDA, Health Canada and EMA, published a

statement in November 2020 stating that follow-up for treatment and placebo arms should continue 'for as long as possible after any regulatory approval' and recommended a follow-up period of 'at least 1 year or more from completion of assigned doses.'<sup>22</sup> Despite this, placebo controlled follow-up, originally planned for 2 years in many trials, was eliminated after a few months, when manufacturers began offering vaccine to placebo recipients within weeks of receiving emergency use authorisations. (Debates took place regarding the ethics of denying placebo recipients vaccine, and proposals to redesign the trials as cross-over trials were not taken up).<sup>23</sup> In addition to ensuring the public accessibility of follow-up data from continuing trials, greater transparency is necessary into the ongoing deliberations and thinking of regulators who are currently evaluating applications from sponsors seeking to move from emergency use authorisations to actual approval or licensure.

Other regulatory documents produced by regulators that can provide greater insight into trials and the evidence development programme include scientific review memos and public assessment reports. In the USA, documents such as review memos and presentations presented to the federal advisory committee are also

released. These are all invaluable, providing insight into regulatory decision making (table 3). At around 50–150 pages, they can be substantially longer than journal articles but still one or two orders of magnitude shorter than clinical study reports, and represent the regulators' analyses of data, not the data itself. Therefore, they should be regarded as complementary to, not substitutes for, trial data.<sup>24</sup>

### Real-time transparency

Transparency should begin as trials get underway, and not be left as a bureaucratic exercise to be conducted after results are announced and decisions are made. This should be the case for all trials, but especially COVID-19 vaccines given their global significance.

Before trialists begin recruiting participants, there are already a variety of pre-study documents in place, and release of these documents would allow for broader awareness and scrutiny of the trials. Are the right endpoints being studied? Are the right populations being recruited? Do informed consent forms convey sufficient information about study purpose?<sup>25</sup> The power of real-time release of trial protocols was on display last summer when some manufacturers—Pfizer, Moderna, Janssen and AstraZeneca (for its US trial)—released study protocols for their phase 3 trials.<sup>26</sup> That transparency not only revealed the inadequacy of what had been disclosed about the studies' primary endpoint in trial registry entries, but helped stimulate public discussion and debate about what primary endpoint was acceptable, including at the FDA's advisory committee the following month—all while the trials were ongoing.<sup>19 27 28</sup>

### Mechanisms to stimulate greater transparency

Although trade-offs will need to be weighed in the context of a global public emergency, there are several immediate steps that governments agencies, regulators, pharmaceutical companies and professional bodies can take to achieve greater transparency of COVID-19 vaccine trials. To start, government agencies and regulators can create infrastructure to support submission and deposits of protocols, informed consent forms, committee charters and other pre-study documents. This could be done by using existing resources such as ClinicalTrials.gov. Moreover, regulatory documents, memos, and scientific reviews used to make decisions on vaccine approval can also be shared on (or linked from) the trial registry entry. Platforms for sharing individual participant data already exist and can be leveraged (eg, Vivli, ClinicalStudyDataRequest.com and YODA), reducing overall monetary cost associated with publishing of trial data.

Drug sponsors and companies can likewise create a dedicated section of their website for pre-study documentation (as some have done). This should not cause an undue burden as such documents are already written and shared with regulators, and they contain no identifying patient information. All that needs to occur is the act of publicly posting a copy. Additionally, sponsors can provide clear and updated statements regarding their timeline for access to electronic individual participant data.

Professional bodies and academic organisations can also play a large role in promoting transparency. Public statements that convey the unacceptability of promises to begin sharing months and years from now, despite vaccine roll-out, would help. More powerfully, professionals could pledge not to endorse new therapeutic agents until full access to data is provided. Doing so would send a strong message that transparency is not a 'nice to have,' but a fundamental component of any intervention that purports to be based on science.<sup>29</sup> These mechanisms and levers can serve

as initial starting points to create greater transparency. Longer-term solutions to advance data transparency may require further changes in policy and law.

### Conclusion

Although progress has been made over the past decades in clinical trial transparency, and there are some successes for COVID-19 vaccines, there is still much room for improvement. The lack of adequate transparency about COVID-19 vaccine trials and their regulation cannot be dismissed as unfortunate, stubborn problems emblematic of the present culture in biomedicine. In a time of increasing public scrutiny, transparency of regulatory decision making leading to the approval of drug treatments and vaccines for COVID-19 is important to ensure patient and stakeholder trust. It is a scientific, moral and ethical imperative that access to complete trial data of these global public health interventions is urgently granted to patients, researchers and other key stakeholders.

Twitter Anisa Rowhani-Farid @AnisaFarid and Peter Doshi @RIATinitiative

**Contributors** ST and PD conceptualised the article and ST wrote the first draft. AR-F and KH led the data extraction and analysis. All authors (ST, AR-F, KH, TJ, PD) were involved in reviewing and editing the final manuscript.

**Funding** The Laura and John Arnold Foundation funds the RIAT Support Center, which supports the salaries of ST, AR-F, KH, TJ and PD.

**Disclaimer** The contents are those of the authors and do not necessarily represent the official views of, nor an endorsement by, FDA/HHS, The University of Maryland, or the US Government.

**Competing interests** KH was supported by the Food and Drug Administration (FDA) of the US Department of Health and Human Services (HHS) as part of a financial assistance award U01FD005946, unrelated to this manuscript, totaling US\$5000 with 100% funded by FDA/HHS. PD has received travel funds from the European Respiratory Society (2012) and Uppsala Monitoring Center (2018); grants from the FDA (through University of Maryland M-CERSI; 2020), Laura and John Arnold Foundation (2017–2022), American Association of Colleges of Pharmacy (2015), Patient-Centered Outcomes Research Institute (2014–2016), Cochrane Methods Innovations Fund (2016–2018) and UK National Institute for Health Research (2011–2014); was an unpaid IMEDS steering committee member at the Reagan-Udall Foundation for the FDA (2016–2020), and is an editor at The BMJ. The Laura and John Arnold Foundation funds the RIAT Support Center, which supports the salaries of ST, AR-F, KH, TJ, and PD. TJ's competing interests are online (<https://restoringtrials.org/competing-interests-tom-jefferson/>).

**Patient consent for publication** Not required.

**Provenance and peer review** Commissioned; externally peer reviewed.

This article is made freely available for personal use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

## ORCID iDs

Sarah Tanveer <http://orcid.org/0000-0002-6408-9855>

Anisa Rowhani-Farid <http://orcid.org/0000-0003-3637-2423>

Tom Jefferson <http://orcid.org/0000-0002-4778-2949>

Peter Doshi <http://orcid.org/0000-0002-7804-4113>

## References

- Turner EH. How to access and process FDA drug approval packages for use in research. *BMJ* 2013;347:f5992.
- European Medicines Agency. European Medicines Agency policy on access to documents (related to medicinal products for human and veterinary use) POLICY/0043 [Internet], 2018. Available: [https://www.ema.europa.eu/en/documents/other/policy/0043-european-medicines-agency-policy-access-documents\\_en.pdf](https://www.ema.europa.eu/en/documents/other/policy/0043-european-medicines-agency-policy-access-documents_en.pdf)
- European Medicines Agency. European Medicines Agency policy on publication of clinical data for medicinal products for human use (policy/0070; ema/240810/2013); 2014.
- Health Canada. Public Release of Clinical Information: consultation [Internet], 2017. Available: <https://www.canada.ca/en/health-canada/programs/consultation-public-release-clinical-information-drug-submissions-medical-device-applications.html>
- Institutions offering data access [Internet] Restoring Invisible and Abandoned Trials (RIAT) Support Center. Available: <https://restoringtrials.org/institutions-offering-data-access>
- EFPIA and PhRMA release joint principles for responsible clinical trial data sharing to benefit patients [Internet]. Available: <https://www.efpia.eu/news-events/the-efpia-view/statements-press-releases/130724-efpia-and-pharma-release-joint-principles-for-responsible-clinical-trial-data-sharing-to-benefit-patients>
- Institute of Medicine. Sharing Clinical Trial Data: Maximizing Benefits, Minimizing Risk [Internet], 2015. Available: <https://www.nap.edu/catalog/18998/sharing-clinical-trial-data-maximizing-benefits-minimizing-risk>
- Doshi P, Jones M, Jefferson T. Rethinking credible evidence synthesis. *BMJ* 2012;344:d7898.
- Eichler H-G, Rasi G. Clinical trial publications: a sufficient basis for healthcare decisions? *Eur J Intern Med* 2020;71:13–14.
- Gøtzsche PC. Why we need easy access to all data from all clinical trials and how to accomplish it. *Trials* 2011;12:249.
- Study to Describe the Safety, Tolerability, Immunogenicity, and Efficacy of RNA Vaccine Candidates Against COVID-19 in Healthy Individuals [Internet]. Available: <https://clinicaltrials.gov/ct2/show/NCT04368728>
- For Whose Benefit? Transparency in the development and procurement of COVID-19 vaccines [Internet]. Available: <https://ti-health.org/content/for-whose-benefit-transparency-in-the-development-and-procurement-of-covid-19-vaccines>
- Xia S, Zhang Y, Wang Y, *et al.* Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *Lancet Infect Dis* 2021;21:39–51.
- Health Canada. Available information for VEKLURY - Submission control number 240551 [Internet], 2020. Available: <https://clinical-information.canada.ca/ci-rc/item/240551>
- Government of Canada HC. Remdesivir authorized with conditions for the treatment of patients in Canada with severe COVID-19 symptoms [Internet], 2020. Available: <https://healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2020/73621a-eng.php>
- Fonseca P. Brazil institute says CoronaVac efficacy above 50% but delays full results [Internet], 2020. Reuters. Available: <https://www.reuters.com/business/healthcare-pharmaceuticals/sinovacs-covid-19-vaccine-over-50-effective-brazil-tests-reports-newspaper-folha-2020-12-23>
- Mallapaty S. China is vaccinating a staggering 20 million people a day. *Nature* 2021. doi:10.1038/d41586-021-01545-3. [Epub ahead of print: 09 Jun 2021].
- Rhodes N, Wright T, Rusu V. For Whose Benefit? Transparency in the development and procurement of COVID-19 vaccines [Internet], 2021. Available: <http://ti-health.org/wp-content/uploads/2021/05/For-Whose-Benefit-Transparency-International.pdf>
- Doshi P. Will covid-19 vaccines save lives? Current trials aren't designed to tell us. *BMJ* 2020;371:m4037.
- Mehrotra DV, Janes HE, Fleming TR, *et al.* Clinical endpoints for evaluating efficacy in COVID-19 vaccine trials. *Ann Intern Med* 2021;174:221–8.
- WHO Ad Hoc Expert Group on the Next Steps for Covid-19 Vaccine Evaluation, Krause PR, Fleming TR, *et al.* Placebo-Controlled Trials of Covid-19 Vaccines - Why We Still Need Them. *N Engl J Med* 2021;384:e2.
- International Coalition of Medicines Regulatory Authorities (ICMRA). Statement on continuation of vaccine trials [Internet]. Available: [http://www.icmra.info/drupal/covid-19/statement\\_on\\_continuation\\_of\\_vaccine\\_trials](http://www.icmra.info/drupal/covid-19/statement_on_continuation_of_vaccine_trials)
- Doshi P. Covid-19 vaccines: in the rush for regulatory approval, do we need more data? *BMJ* 2021;373:n1244.
- Jefferson T, Jones MA, Doshi P, *et al.* Neuraminidase inhibitors for preventing and treating influenza in adults and children. *Cochrane Database Syst Rev* 2014;19.
- Doshi P, Hur P, Jones M, *et al.* Informed consent to study purpose in randomized clinical trials of antibiotics, 1991 through 2011. *JAMA Intern Med* 2017;177:1452–9.
- Doshi P. Covid-19 vaccine trial protocols released. *BMJ* 2020;371:m4058.
- Doshi P, Topol E. Opinion, 2020. Available: <https://www.nytimes.com/2020/09/22/opinion/covid-vaccine-coronavirus.html>
- U.S. Food and Drug Administration. 161st vaccines and related biological products Advisory Committee (VRBPAC) meeting, 2020. Available: <https://www.fda.gov/media/143982/download>
- Johnson RM, Doshi P, Healy D. Covid-19: should doctors recommend treatments and vaccines when full data are not publicly available? *BMJ* 2020;370:m3260.
- Center for Drug Evaluation, Research. E3 structure and content of clinical study reports, 2020. Available: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/e3-structure-and-content-clinical-study-reports>
- Doshi P, Dickersin K, Healy D, *et al.* Restoring invisible and abandoned trials: a call for people to publish the findings. *BMJ* 2013;346:f2865.
- Glossary [Internet] Restoring Invisible and Abandoned Trials (RIAT) Support Center. Available: <https://restoringtrials.org/glossary>