

Evidence-Based Medicine

Contents

| | |
|---|-------------------|
| Purpose and Procedure | |
| Additional Articles Abstracted in ACP Journal Club | |
| EBM NOTEBOOK | |
| Where do pretest probabilities come from? | 68 |
| Bringing evidence to the point of care | 70 |
| David Sackett on the move; Paul Glasziou takes over | 71 |
| THERAPEUTICS | |
| Oral methylprednisolone improved symptoms and disability resulting from multiple sclerosis attacks | 73 |
| Interferon- β 1a reduced relapses at 2 years in relapsing-remitting multiple sclerosis | 74 |
| Interferon- β 1b reduced the progression of secondary progressive multiple sclerosis | 75 |
| Xylitol syrup and xylitol chewing gum were both effective in preventing acute otitis media | 76 |
| Review: Amoxicillin and folate inhibitors are as effective as newer, more expensive antibiotics for acute sinusitis | 77 |
| Review: Insecticide-treated bednets and curtains reduce overall mortality and incidence of malaria | 78 |
| 2 or 4 sessions of counselling reduced occurrence of sexually transmitted diseases | 79 |
| Review: Measures to control house dust mites are not effective in patients with asthma who are sensitive to mites | 80 |
| Review: Successful <i>Helicobacter pylori</i> eradication prevents ulcer recurrence | 81 |
| Review: St. John's wort is more effective than placebo for treating depressive disorders | 82 |
| Flutamide did not prolong survival and increased toxic effects after orchiectomy in metastatic prostate cancer | 83 |
| Review: Postoperative radiotherapy decreases survival in early-stage non-small-cell lung cancer | 84 |
| Preservation of the intercostobrachial nerve during axillary node clearance for breast cancer reduced sensory deficits | 85 |
| Second-eye cataract surgery was beneficial in otherwise-healthy patients | 86 |
| Review: Carotid endarterectomy modestly reduces ipsilateral stroke in asymptomatic stenosis | 87 |
| Early surgery was more expensive than surveillance for small abdominal aortic aneurysms but slightly improved health perception and bodily pain | 88 |
| Early surgery did not reduce 6-year mortality in patients with small abdominal aortic aneurysms | 89 |
| DIAGNOSIS | |
| D-dimer assay was useful for ruling out suspected pulmonary embolism | 90 |
| PROGNOSIS | |
| Advanced age predicted risk for recurrent stroke after first-ever stroke | 91 |
| ETIOLOGY | |
| Review: Bias may contribute to the association between vasectomy and prostate cancer | 92 |
| Oral contraceptive use was associated with myocardial infarction and ischaemic stroke but not angina | 93 |
| QUALITY IMPROVEMENT | |
| Nurse-run clinics in primary care increased secondary prevention in coronary artery disease | 94 |
| A nurse-run program for congestive heart failure reduced time to readmission | 95 |
| Review: Computer-based clinical decision support systems can improve physician performance in some areas | 96 |
| Glossary | Inside back cover |

Best Evidence 3, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1998, is available to institutions for £95, BMA members for £43, and to non-members for £62. To order, contact BMJ Publishing Group at 0171 383 6270. Information about Best Evidence 3 is also available at <http://www.bmjpg.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.