

Evidence-Based Medicine

Contents

Purpose and Procedure	130	Laparoscopically assisted vaginal hysterectomy resulted in less pain and shorter hospital stay than abdominal hysterectomy	149
Additional Articles Abstracted in ACP Journal Club	132	Review: Support interventions reduce cessation of breast-feeding within 2 months of delivery	150
EBM Notebook		Review: Adjuvant chemotherapy improves recurrence-free intervals and recurrence-free survival in soft-tissue sarcoma	151
Searching for evidence: principles and practice	133	Laparoscopic appendectomy was better than open appendectomy for subjective full recovery but not for duration of sick leave	152
THERAPEUTICS		Zidovudine reduced vertical transmission of HIV infection in breast-fed infants	153
Early surgery was better than watchful waiting for language development in children with persistent otitis media with effusion	137	A pacemaker reduced the recurrence of syncope in patients with severe vasovagal syncope	154
Review: Magnesium sulfate is effective for severe acute asthma treated in the emergency department	138	DIAGNOSIS	
A structured discharge package given by a nurse reduced hospital readmission and reattendance in children with asthma	139	Review: Physical signs are not useful for detecting non-blood-loss hypovolemia	155
Influenza vaccine prevented influenza infection in health care workers	140	Review: Echocardiography, SPECT, and immediate angiography are cost-effective for detecting CAD	156
Exercise training reduced mortality and cardiac events and improved quality of life in chronic heart failure	141	Albumin-to-creatinine ratio in a timed overnight urine sample was accurate for screening for microalbuminuria in diabetes mellitus	157
Review: 6 clinical conditions can be modified to reduce risk for a first stroke	142	ETIOLOGY	
Review: Antihypertensive drugs reduce stroke in patients 80 years of age or older	144	Evidence of greater medical knowledge was associated with increased malpractice claims	158
Review: Medication and cognitive behaviour therapy control symptoms of bulimia nervosa	145	Symptoms in patients with syncope did not predict death or recurrence	159
Cognitive behavior therapy, temazepam, or both improved short-term outcomes for older adults with persistent insomnia	146	Higher hospital volume was associated with lower operative mortality and shorter length of stay after hepatic resection	160
Intervention by paraprofessional advocates helped abused women meet their needs and goals and reduced violence	147	Glossary	Inside back cover
10 mg of mifepristone prevented pregnancy as effectively as 600 mg within 5 days of unprotected intercourse	148		

Best Evidence 3, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1998, is available to institutions for £95, BMA members for £43, and to non-members for £62. To order, contact BMJ Publishing Group at 0171 383 6270. Information about Best Evidence 3 is also available at <http://www.bmjpg.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.