

Evidence-Based Medicine

Contents

Purpose and procedure	2	Glucocorticoids reduced short-term treatment failure in exacerbations of chronic obstructive pulmonary disease	18
Additional articles abstracted in ACP journal club	3	A 14-day regimen of high-dose inhaled corticosteroids was more effective than placebo for persistent nocturnal cough ..	19
EBM notebook		Buccal midazolam may be a useful alternative to rectal diazepam for treating acute seizures in refractory epilepsy ...	20
Advances in evidence-based information resources for clinical practice	4	Pergolide reduced involuntary periodic leg movements and improved sleep in the restless legs syndrome	21
Appendix	5	Review: lower doses of antidepressant drugs are effective and have fewer adverse effects in depression	22
Journals reviewed for this issue	6	Review: studies with sufficient follow-up do not show a clear benefit for pharmacotherapy in alcohol dependence	23
Resource corner	7	Transcervical resection of the endometrium was effective and acceptable at up to 2 years for heavy menstrual bleeding	24
Therapeutics		Laparoscopic hernia repair reduced recovery time and days of postoperative analgesia use better than Shouldice repair ..	25
Review: aspirin reduces the risk for stroke in patients with previous transient ischaemic attack or stroke but does not have a dose-response effect	9	Review: avulsion with phenol prevents recurrence of ingrown toenails	26
Review: low-dose thiazides are the most effective first-line drugs for hypertension	10	The <i>Baby Check</i> booklet did not affect frequency of health service use in infants ≤ 6 months of age	27
Spironolactone reduced mortality in severe congestive heart failure	11	Diagnosis	
Review: long-term β -blocker use reduces mortality after myocardial infarction	12	Review: the physical examination can exclude the diagnosis of meningitis in low-risk adults	28
n-3 polyunsaturated fatty acids reduced mortality and morbidity after recent myocardial infarction	13	Review: sensitive thyrotropin testing in unselected inpatients has low diagnostic accuracy	29
Atorvastatin was at least as effective as PTCA for reducing ischaemic events in patients with stable coronary artery disease	14	Whole-blood antibody tests were not highly sensitive for detecting <i>Helicobacter pylori</i> infection	30
Review: acellular vaccines prevent pertussis as effectively as whole-cell vaccines with fewer adverse events	15	The ^{13}C -urea blood test is accurate for detecting <i>Helicobacter pylori</i> infection	31
Review: β -agonists plus ipratropium bromide improve airflow obstruction in adults with acute asthma	16	Aetiology	
Inhaled budesonide plus oral prednisone reduced relapse after discharge from the emergency department for acute asthma	17	Diastolic blood pressure and weight gain predicted future risk for hypertension	32
		Glossary	Inside back cover

Best Evidence 3, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1998, is available to institutions for £95, BMA members and £43, and to non-members for £62. To order, contact BMJ Publishing Group at 0171 383 6270. Information about Best Evidence 3 is also available at <http://www.bmjpub.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.