The ¹³C-urea blood test is accurate for detecting *Helicobacter pylori* infection

Chey WD, Murthy U, Toskes P, et al. *The ¹³C-urea blood test accurately detects active Helicobacter pylori infection: a United States, multicenter trial.* Am J Gastroenterol. 1999 Jun;94:1522-4.

QUESTION: How accurate is the ¹³C-urea blood test for detecting *Helicobacter pylori* infection?

Design

Blinded comparison of the ¹³C-urea blood test with tests based on endoscopic biopsy.

Setting

5 centres in the United States (Ann Arbor, Michigan; Syracuse, New York; Gainesville, Florida; Savannah, Georgia; and Los Angeles, California).

Patients

121 patients (mean age 49 y, 51% men) who were referred for endoscopy. Exclusion criteria included treatment for *H. pylori* infection in the previous year or use of antibiotics or bismuth in the previous month or proton-pump inhibitors in the previous 7 days.

Description of tests and diagnostic standards

Patients received ¹³C-urea, 125 mg dissolved in 75 ml of water. 30 minutes later, a 3-ml blood sample was obtained by venipuncture and analysed by gas isotope ratio mass spectrometry. The 3 diagnostic standards were histological evidence of *H. pylori* infection in biopsies obtained from the body and antrum of the stomach, a positive result for both histological and rapid urease testing (RUT) (patients with discordant histological and

RUT results were considered uninfected), and a positive result for either histological testing or RUT.

Main outcome measures

Sensitivity and specificity for detecting *H. pylori* infection.

Main results

The table shows sensitivities, specificities, and likelihood ratios. Results for the ¹³C-urea blood test did not differ from those for RUT ($p > 0.2$).

Conclusion

The ¹³C-urea blood test was similar to rapid urease testing and had high sensitivity and specificity for detecting *Helicobacter pylori* infection.

*Test characteristics for detecting Helicobacter pylori infection**

Diagnostic standards	Tests	Sensitivity (95% CI)	Specificity (CI)	+LR	-LR
Histological testing	¹³ C-UBT	89% (85 to 93)	96% (94 to 98)	19.9	0.1
	RUT	87% (75 to 95)†	96% (87 to 99)†	19.4†	0.1†
Histological testing and RUT	¹³ C-UBT	94% (87 to 100)‡	91% (85 to 97)‡	10.4	0.1
Histological testing or RUT	¹³ C-UBT	88% (80 to 96)‡	98% (95 to 100)‡	44.0	0.1

*¹³C-UBT = ¹³C-urea blood test; RUT = rapid urease testing. LR = defined in glossary and calculated from data in article.

†Calculated from data supplied by author.

‡CIs provided by author.

COMMENTARY

(continued from page 30)

The combination of reference standard error, spectrum bias, and a greater potential for operator error means that caution should be used when extrapolating these results to the office setting.³

Unfortunately, although the absolute values of the performance of the ¹³C-urea blood test are greater than the whole-blood antibody tests, the confidence intervals overlap, which means that we cannot be certain that the difference is robust. In any case, clinical differences between the 2 types of tests will be small because, at most, only 20% of patients will benefit from the "test and eradicate" strategy,⁴ and the absolute difference in sensitivity of the tests is only 5% to 10%.¹ Only 2 patients in 100 might be missed with the antibody test. An evaluation of the tests in the office setting with larger samples and a health economic analysis are needed before an informed choice can be made between whole-blood tests and ¹³C-urea-based tests for applying the "test and eradicate" strategy in the office.

Brendan Delaney, MD, BmBCh
University of Birmingham
Birmingham, UK

Source of funding: not stated.

For correspondence: Dr WD Chey, University of Michigan Medical Center, 3912 Taubman Center, Box 0362, Ann Arbor, MI 48109, USA. FAX 734-936-7392.

- Calam J. Clinicians' guide to *Helicobacter pylori*. London: Chapman and Hall, 1996.
- Heaney A, Collins JS, Watson RG, et al. A prospective randomised trial of a "test and treat" policy versus endoscopy based management in young *Helicobacter pylori* positive patients with ulcer-like dyspepsia, referred to a hospital clinic. Gut 1999;45:186-90.
- Irwig L, Tosteson AN, Gatsonis C, et al. Guidelines for meta-analyses evaluating diagnostic tests. Ann Intern Med 1994;120:667-76.
- McColl K, Murray L, El-Omar E, et al. Symptomatic benefit from eradicating *Helicobacter pylori* infection in patients with non-ulcer dyspepsia. N Engl J Med 1998;339:1869-74.