

Evidence-Based Medicine

Contents

Purpose and procedure	34	More people with depression continued treatment with fluoxetine than with desipramine or imipramine.....	51
Additional articles abstracted in ACP journal club	35	Methylprednisolone injection improved symptoms for 1 year in patients with the carpal tunnel syndrome	52
EBM notebook		Review: exercise therapy may reduce pain and disability in osteoarthritis of the hip or knee	53
Assessing allocation concealment and blinding in randomised controlled trials: why bother?	36	Raloxifene reduced vertebral fractures in postmenopausal women	54
Incorporating allocation concealment and blinding in randomised controlled trials	38	Trimethoprim sulfamethoxazole decreased morbidity and mortality in HIV-1 infected patients with tuberculosis.....	55
Corrections	38	Cutaneous lumbosacral injections of sterile water were more effective than “placebo” injections for relieving first stage labour pain	56
Resource corner	39	Tamoxifen added to lumpectomy and radiation therapy reduced breast cancer events in ductal carcinoma in situ	57
Therapeutics		Use of an impervious wound edge protector decreased postoperative wound infection	58
Review: glucocorticoids improve symptoms of croup within 6 hours	41	Omeprazole was better than cisapride or placebo for controlling heartburn at 4 and 8 weeks	59
Review: antibiotics have a slight beneficial effect on acute bronchitis.....	42	Review: health care decision aids improve knowledge, decrease decisional conflict, and increase active participation ..	60
Review: penicillin V or amoxicillin is better than placebo and equal to non-penicillins for acute maxillary sinusitis	43	Diagnosis	
Intranasal, live, attenuated influenza virus vaccine reduced severe febrile illness in healthy adults	44	Review: screening tests are not as accurate as overnight polysomnography for the diagnosis of adult sleep apnoea ...	61
Attendance at an asthma clinic did not reduce asthma morbidity more than did standard medical treatment.....	45	Aetiology	
Grass pollen immunotherapy for 3 to 4 years was still effective 3 years after being discontinued	46	Surgical volume was not related to 30 day mortality in 8 common operations	62
Ramipril reduced mortality and cardiovascular morbidity in high risk adults	47	Review: high levels of homocysteine are associated with an increased risk for cardiovascular disease	63
Review: multiple interventions modestly reduce cardiovascular risk factors in primary prevention, but effects on mortality are uncertain	48	Quality improvement	
Gemfibrozil reduced the risk for coronary events in men with coronary artery disease and low levels of high density lipoprotein cholesterol.....	49	Review: interactive, but not didactic, continuing medical education is effective in changing physician performance	64
Empirical eradication therapy was better than endoscopy based management for <i>Helicobacter pylori</i> infection and dyspepsia	50	Glossary	Inside back cover

Best Evidence 3, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1998, is available to institutions for £95, BMA members for £43, and to non-members for £62. To order, contact BMJ Publishing Group at +44 (0)20 7383 6270. Information about Best Evidence 3 is also available at <http://www.bmj.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.