

Evidence-Based Medicine

Contents

Purpose and procedure	66	Conventional and newer antihypertensive drugs had similar efficacy in elderly people with hypertension	85
Additional articles abstracted in ACP journal club	67	Paroxetine was effective for reducing symptoms in social phobia	86
EBM notebook		Antidepressants plus cognitive therapy reduced relapse in residual depression	87
Identifying systematic reviews: key resources	68	Review: lithium augmentation increases treatment response in refractory depression	88
Resource corner	70	Women with breast cancer were more satisfied with general practitioner care than with outpatient clinic care	89
Therapeutics		Review: β sitosterols improve urinary symptoms in the short term in men with benign prostatic hyperplasia	90
Hyaluronic acid improved walking ability and short term knee function in osteoarthritis of the knee	73	Daily use of β carotene supplements did not prevent skin cancer, but daily sunscreen use reduced the incidence of squamous cell carcinomas	91
Rofecoxib caused fewer endoscopic gastroduodenal ulcers than ibuprofen in osteoarthritis	73	Diagnosis	
Celecoxib was similar to naproxen for rheumatoid arthritis with fewer endoscopic ulcers	75	A negative D-dimer result and low risk clinical status effectively ruled out DVT in symptomatic patients	92
Review: early mobilisation may be better than bed rest for medical conditions and after surgery	76	A negative D-dimer test result alone or combined with low risk clinical status effectively ruled out symptomatic DVT	93
An 8 session exercise programme was effective for subacute or chronic low back pain	77	A fasting plasma glucose level ≥ 7.8 mmol/l was sensitive but not specific for identifying poor glycaemic control	94
Review: evidence supports surgery for lumbar disc prolapse but is insufficient for degenerative lumbar spondylosis	78	Aetiology	
Spacers made from sealed cold-drink bottles were as effective as conventional spacers in children with acute asthma	79	HIV transmission risk during breast feeding was greatest during the early months of life	95
Zanamivir speeds symptom relief in influenza	80	Clinical prediction guide	
Oseltamivir once or twice daily safely prevented influenza	81	The high risk criteria of a clinical prediction model were specific but not sensitive for predicting ectopic pregnancy	96
Review: antithrombotic agents prevent stroke in non-valvular atrial fibrillation	82	Glossary	Inside back cover
Review: psychoeducational programmes reduce MI recurrence and improve some physical health outcomes	83		
A multidisciplinary, home based intervention reduced deaths and readmissions in patients with chronic congestive heart failure	84		

Best Evidence 4, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1999, is available to institutions for £103, BMA members for £50, and to non-members for £70. To order, contact BMJ Publishing Group at +44 (0)20 7383 6270. Information about Best Evidence 4 is also available at <http://www.bmjpg.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.