

# Evidence-Based Medicine

## Contents

<b>Purpose and procedure</b> .....	98	Review: antidepressive agents improve symptoms and reduce pain in patients with functional gastrointestinal disorders....	115
<b>Additional articles abstracted in ACP Journal Club</b> .....	99	<i>Hypericum</i> extract was better than placebo and equivalent to imipramine for moderate depression.....	116
<b>EBM notebook</b>		Review: inadequate data exist on prevention and treatment strategies for foot ulcer in diabetes mellitus.....	117
The need and means for evidence-based medicine in developing countries.....	100	Review: antimicrobial prophylaxis reduces asymptomatic bacteriuria in patients with neurogenic bladder.....	118
<b>Resource corner</b> .....	102	Alendronate reduced days of bed rest and limited activity in postmenopausal women with osteoporosis and existing fractures .....	119
<b>Correction</b> .....	104	Local injection, naproxen, and simple analgesia led to similar 1 year rates of symptom relief in lateral epicondylitis .....	120
<b>Therapeutics</b>		Review: injection treatment is not better than placebo for relieving pain in benign chronic low back pain .....	121
Anti-androgen treatment immediately after surgery for prostate cancer improved survival .....	105	Pacifiers have greater analgesic effect than sweet solutions during venipuncture in full term newborns.....	122
Review: providing recordings or summaries of consultations may help patients with cancer .....	106	Review: the extended tip spatula is more effective than the Ayre's spatula for detecting karyosis in cervical smears.....	123
Review: ipratropium bromide with $\beta$ -agonists improves pulmonary function and reduces admissions to hospital in acute asthma.....	107	<b>Diagnosis</b>	
Review: oral <i>Haemophilus influenzae</i> vaccination reduces the number and severity of recurrent bronchitis episodes.....	108	Review: newer technologies improve sensitivity in detecting uterine cervical lesions but at a substantially increased cost .	124
Oral montelukast was better than inhaled salmeterol for reducing exercise induced bronchoconstriction in adults with asthma.....	109	Review: diagnostic tests for bacterial rhinosinusitis have moderate accuracy; antibiotics reduce clinical failures.....	125
High dose lisinopril was more effective than low dose for reducing combined mortality and cardiovascular events in congestive heart failure .....	110	The Uriscreeen test was not better than standard urinalysis and dipstick tests for detecting urinary tract infection in children .....	126
Review: high dose and moderate dose oral anticoagulants reduce events in coronary artery disease but increase major bleeding and are no more effective than aspirin .....	111	The near-patient test FlexSure had low sensitivity and high specificity for detecting <i>Helicobacter pylori</i> infection.....	127
Low volume, high frequency cows' milk feedings sped recovery from diarrhoea in infants .....	112	<b>Aetiology</b>	
<i>Lactobacillus GG</i> reduced diarrhoea incidence in children treated with antibiotics .....	113	Older adults who reported strain when caring for a spouse with disabilities had increased mortality .....	128
Review: newer and older antidepressants have similar efficacy and total discontinuation rates but different side effects.....	114	<b>Glossary</b> .....	Inside back cover

**Best Evidence 4**, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1999, is available to institutions for £103, BMA members for £50, and to non-members for £70. To order, contact BMJ Publishing Group at +44 (0)20 7383 6270. Information about Best Evidence 4 is also available at <http://www.bmj.com/data/ebm.htm>.

*Evidence-Based Medicine* is published bimonthly with the support of the NHS Executive Research and Development Programme.