

Evidence-Based Medicine

Contents

| | | | |
|---|-----|--|-------------------|
| Purpose and procedure | 98 | Review: antidepressive agents improve symptoms and reduce pain in patients with functional gastrointestinal disorders.... | 115 |
| Additional articles abstracted in ACP Journal Club | 99 | <i>Hypericum</i> extract was better than placebo and equivalent to imipramine for moderate depression..... | 116 |
| EBM notebook | | Review: inadequate data exist on prevention and treatment strategies for foot ulcer in diabetes mellitus..... | 117 |
| The need and means for evidence-based medicine in developing countries..... | 100 | Review: antimicrobial prophylaxis reduces asymptomatic bacteriuria in patients with neurogenic bladder..... | 118 |
| Resource corner | 102 | Alendronate reduced days of bed rest and limited activity in postmenopausal women with osteoporosis and existing fractures | 119 |
| Correction | 104 | Local injection, naproxen, and simple analgesia led to similar 1 year rates of symptom relief in lateral epicondylitis | 120 |
| Therapeutics | | Review: injection treatment is not better than placebo for relieving pain in benign chronic low back pain | 121 |
| Anti-androgen treatment immediately after surgery for prostate cancer improved survival | 105 | Pacifiers have greater analgesic effect than sweet solutions during venipuncture in full term newborns..... | 122 |
| Review: providing recordings or summaries of consultations may help patients with cancer | 106 | Review: the extended tip spatula is more effective than the Ayre's spatula for detecting karyosis in cervical smears..... | 123 |
| Review: ipratropium bromide with β -agonists improves pulmonary function and reduces admissions to hospital in acute asthma..... | 107 | Diagnosis | |
| Review: oral <i>Haemophilus influenzae</i> vaccination reduces the number and severity of recurrent bronchitis episodes..... | 108 | Review: newer technologies improve sensitivity in detecting uterine cervical lesions but at a substantially increased cost . | 124 |
| Oral montelukast was better than inhaled salmeterol for reducing exercise induced bronchoconstriction in adults with asthma..... | 109 | Review: diagnostic tests for bacterial rhinosinusitis have moderate accuracy; antibiotics reduce clinical failures..... | 125 |
| High dose lisinopril was more effective than low dose for reducing combined mortality and cardiovascular events in congestive heart failure | 110 | The Uriscreeen test was not better than standard urinalysis and dipstick tests for detecting urinary tract infection in children | 126 |
| Review: high dose and moderate dose oral anticoagulants reduce events in coronary artery disease but increase major bleeding and are no more effective than aspirin | 111 | The near-patient test FlexSure had low sensitivity and high specificity for detecting <i>Helicobacter pylori</i> infection..... | 127 |
| Low volume, high frequency cows' milk feedings sped recovery from diarrhoea in infants | 112 | Aetiology | |
| <i>Lactobacillus GG</i> reduced diarrhoea incidence in children treated with antibiotics | 113 | Older adults who reported strain when caring for a spouse with disabilities had increased mortality | 128 |
| Review: newer and older antidepressants have similar efficacy and total discontinuation rates but different side effects..... | 114 | Glossary | Inside back cover |

Best Evidence 4, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1999, is available to institutions for £103, BMA members for £50, and to non-members for £70. To order, contact BMJ Publishing Group at +44 (0)20 7383 6270. Information about Best Evidence 4 is also available at <http://www.bmjpub.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.