

Evidence-Based Medicine

Contents

| | | | |
|--|-----|--|-------------------|
| Purpose and procedure | 130 | Amoxicillin resolved otitis media symptoms in young children better than did placebo at day 4 but not at day 11..... | 149 |
| Additional articles abstracted in <i>ACP Journal Club</i> | 131 | Cisapride was effective in the short term for constipation in children | 150 |
| EBM Notebook | | Routine testing before cataract surgery did not reduce medical adverse events | 151 |
| Clinical assessment of the reliability of the examination (CARE) | 132 | Diagnosis | |
| Resource corner | 135 | Tachypnea is a useful predictor of pneumonia in children with acute respiratory infection..... | 152 |
| Therapeutics | | A respiratory rate of ≥ 60 breaths per minute had high sensitivity for detecting hypoxia in infants | 153 |
| Vitamin E did not reduce myocardial infarction, death, or stroke in adults at high risk for cardiovascular events..... | 137 | Biliary scintiscan had high sensitivity and specificity for predicting pathological findings in the common bile duct... | 154 |
| Review: isolated systolic hypertension increases mortality and morbidity in elderly people and should be treated..... | 138 | Aetiology | |
| Inhaled budesonide and prednisone led to similar rates of asthma relapse after emergency department discharge | 139 | Patients with carotid artery stenosis had a high risk for stroke, and some types of stroke were not prevented by endarterectomy | 155 |
| An outpatient multidisciplinary pulmonary rehabilitation programme was effective in disabling chronic lung disease.. | 140 | Quality improvement | |
| Review: treatment induced blood pressure reductions in pregnancy may be associated with decreased fetal growth... | 141 | A critical pathway reduced resource use without compromising safety and effectiveness in community acquired pneumonia..... | 156 |
| Controlled and extended release metoprolol reduced death, hospital admissions, and symptoms in chronic heart failure. | 142 | Feedback to physicians plus telephone care management improved outcomes in primary care patients with depression..... | 157 |
| Review: antidepressants are effective for clinical improvement in unexplained physical symptoms and syndromes | 143 | Practice based education did not increase recognition of depression by primary care physicians nor improve the outcome of depression | 158 |
| Review: early treatment of acute herpes zoster may prevent or shorten the duration of postherpetic neuralgia | 144 | Economics | |
| Manual physical therapy and exercise improved function in osteoarthritis of the knee..... | 145 | Brief physician advice to problem drinkers reduced alcohol intake and societal costs..... | 159 |
| Review: glucosamine and chondroitin improve outcomes in osteoarthritis, but the magnitude of effect is unclear | 146 | Clinical prediction guide | |
| Review: evidence is unclear that preventive home visits for elderly people in the community improve mortality or health outcomes | 147 | A clinical prediction tool that includes modifiable risk factors predicted functional decline in elderly women..... | 160 |
| Review: long-term effectiveness of non-drug treatments for sleep problems in young children is more evident | 148 | Glossary | Inside back cover |

Best Evidence 4, a complete electronic archive of *ACP Journal Club* and *Evidence-Based Medicine* through 1999, is available to institutions for £103, BMA members and £50, and to non-members for £70. To order, contact BMJ Publishing Group at +44(0)20 7383 6270. Information about Best Evidence 4 is also available at <http://www.bmjpub.com/data/ebm.htm>.

Evidence-Based Medicine is published bimonthly with the support of the NHS Executive Research and Development Programme.