


Evidence-Based Medicine

Editors: Brian Haynes & Paul Glasziou



Contents

Purpose and procedure	66	Budesonide and nedocromil did not improve lung function in children with asthma	84
Additional articles abstracted in <i>ACP Journal Club</i>	67	Review: lipid lowering drugs decrease coronary artery disease (CAD) events but not all cause or CAD mortality in men with no history of CAD	85
EBM notebook: A graphic framework for teaching critical appraisal of randomised controlled trials	68	Review: falls can be prevented in older people, but interventions should be multifaceted and targeted	86
Resource corner	70	Racecadotril was effective for severe watery diarrhoea in children	87
Therapeutics		Review: misoprostol, double dose H ₂ receptor antagonists, and proton pump inhibitors reduce GI ulcers in long term NSAID use	88
Planned caesarean section had lower risk for perinatal morbidity and mortality than planned vaginal birth in breech presentation	73	Faecal occult blood screening reduced the incidence of colorectal cancer	89
A midwife led debriefing session after operative childbirth did not reduce postpartum depression	74	Fibre supplementation increased the risk for recurrent adenomas, and calcium supplementation did not prevent recurrence	90
Review: selective serotonin reuptake inhibitors reduce symptoms in premenstrual syndrome	75	Contamination of urine specimens did not differ with collection technique in women with acute dysuria	91
An active exercise and posture protocol reduced pain in acute whiplash injuries	76	Antibiotics as an adjunct to phenol matrixectomy did not decrease healing time of ingrown toenails	92
Review: antidepressants improve symptoms of fibromyalgia	77	Bug busting (wet combing) was less effective than malathion lotion for eliminating head lice in children	93
Interferon β -1a prevented the development of clinically definite multiple sclerosis after a first demyelinating event	78	Diagnosis	
Glyburide was as safe and effective as insulin in gestational diabetes	79	Review: hand symptom diagrams, weak thumb abduction, and hypalgesia are helpful in diagnosing carpal tunnel syndrome	94
Targeting postprandial blood glucose concentrations reduced haemoglobin A _{1c} concentrations in type 2 diabetes mellitus	80	Dual section helical computed tomography had high sensitivity and specificity for detecting acute pulmonary embolism	95
Review: self monitoring does not improve blood glucose control in type 2 diabetes mellitus	81	Clinical prediction guide	
Review: some non-antibiotic treatments are effective for relieving acute sore throat	82	A 2 factor model helped to rule out early stage necrotising fasciitis	96
Influenza vaccination of healthy adults was effective during the 1998–9 flu season but not the 1997–8 season	83		